


# 研究彈薪表單填寫教學

(A類-獎勵、N類-新聘、Y類-延攬)

研發處113.3.20修正

本表僅作為輔助分辨可適用之申請類型，補助與否仍需以視申請人資格及實際績效而定。



# N類-新聘

【辦理期程:每年二梯次(2月、6月)】

申請表(首次申請者)

申請表(續撥獎勵者)

# 1. 首次申請者

必備條件:

- ▶ 本校新聘任期三年(含)以下。
- ▶ 執行國科會專題研究計畫之教學研究人員。
- ▶ 以國內第一次延攬聘任者為限，不含由國內公私立大專校院或學術研究機關(構)延攬之人員。
- ▶ 需特別留意是否已申請/聘任其他競合性彈性薪資(例：資安彈薪)。

## 2. 續撥申請者

必備條件:

- ▶ 第一年至少發表**一篇**收錄於 Scopus 或 WOS 資料庫之國際學術期刊論文或國際研討會論文。
- ▶ 第二年及第三年至少需各發表收錄**一篇**和**二篇**於 Scopus 或 WOS 資料庫之國際學術期刊論文(不含國際研討會論文)。
- ▶ 獎勵期限每人至多三年，且不得中斷聘期，並**須每年提出申請**，經審查通過後，由本校依當年度可使用經費總額核予獎勵金，按月撥付。

# Y類-延攬

【辦理期程:每年一次(6月底申請)】

申請表

## 注意事項:

- ▶ 可採計教師於前一任職單位近五年傑出績效表現。
- ▶ 近五(學)年曾執行政府計畫三個(學)年度以上。
- ▶ 申請時務必留意須將預期績效分年度並具體列出以供審查與考核。
- ▶ 延攬(Y)與新聘(N)申請條件不相同，不可同時申請。
- ▶ 延攬(Y)需與獎勵(A)之第八、九級申請人一同進行排序(例：Y類申請人等同A類第X名申請者)

# A類-獎勵

【辦理期程:每年二梯次(6月底申請)】

申請表



## 注意事項:

- ▶ 採計近五年傑出績效表現。
- ▶ 以本校名義發表論文、執行國科會計畫、產學合作計畫、技術移轉等項目綜合考評機制。
- ▶ 共分為九個等級，依等級核發獎勵金，可勾多個級別，擇優辦理。
- ▶ 第5~7級的申請名單不需經學院排名。
- ▶ 第8、9級的申請需先經學院排名後提供排序名單，再由學校依各學院貢獻度依比例排名。

## 獎項方面

- 需提供佐證資料
- 前一年度且累積二次以上  
EX: 113年度申請・110年度獲獎+108年度曾獲獎即有符合

申請項目(請勾選並檢附-表 A「近五年內之傑出績效說明表」) ※本次補助起始日:113.8.1	
第一級 至 第四級	申請級數：第_____級 第_____項 適用標準：_____ ※ 請依「附表一：本校獎勵特殊優秀研究人才支給標準表」之等級適用標準自行填寫
第五級	<input type="checkbox"/> 1.國際知名之國家院士 <input type="checkbox"/> 2.曾(現)任國際著名大學之講座 <input type="checkbox"/> 3.前一年度以本校名義獲國科會傑出研究獎且累積獲獎次數達二次以上 <input type="checkbox"/> 4.當學年度獲聘為本校終身講座教授者 <input type="checkbox"/> 5.前一年度以本校名義發表之論文點數達250點以上(不含研討會論文)
第六級	<input type="checkbox"/> 1.前一年度以本校名義獲國科會傑出研究獎 <input type="checkbox"/> 2.當學年度獲聘為本校專任講座教授者 <input type="checkbox"/> 3.前一年度以本校名義發表之論文點數達140點以上(不含研討會論文)
第七級	<input type="checkbox"/> 1.當學年度獲聘為本校終身特聘教授者 <input type="checkbox"/> 2.前一年度獲本校傑出研究獎或傑出產學合作獎且累積次數達二次以上者 <input type="checkbox"/> 3.前一年度以本校名義發表之論文點數達80點以上(不含研討會論文) <input type="checkbox"/> 4.前一年度以本校名義所獲得之產學合作計畫點數達240點以上且管理費納入校務基金超過150萬元 <input type="checkbox"/> 5.前一年度以本校名義所獲得之實收技術移轉金點數達175點以上且管理費納入校務基金超過50萬元 ※ 以本級第4、5項申請者，「傑出績效說明表」請先至產學處確認核章後，再送回各系所審查。
第八級 及 第九級	<input type="checkbox"/> 1.當學年度獲聘為本校特聘教授者 <input type="checkbox"/> 2.前一年度獲本校傑出研究獎者 <input type="checkbox"/> 3.前一年度獲本校傑出產學合作獎者 <input type="checkbox"/> 4.基本門檻：近五年內曾主持國科會各型計畫，並滿足下列條件之一者 <ul style="list-style-type: none"> <li><input type="checkbox"/> (1)近五年以本校名義發表之重要學術論著績效點數12點。 設計學院及人社學院教師得採計 TSSCI/THCI 期刊論文；人文、設計、藝術或社會科學領域教師得以學術專書著作或專章申請。</li> <li><input type="checkbox"/> (2)以本校名義主持國科會各類型計畫，五年內之總金額，以五年內之計畫總金額為通過標準，其標準由各學院依相關程序訂定之。</li> <li><input type="checkbox"/> (3)近五年以本校名義所獲得之產學合作計畫累計總金額超過1000萬元(績效點數200點)且管理費納入校務基金超過150萬元者。</li> <li><input type="checkbox"/> (4)近五年以本校名義所獲之實收技術移轉金累計總金額超過250萬元(績效點數125點)且管理費納入校務基金超過50萬元者。</li> </ul>
申請第4項 (1)(2)院推薦 排序	
(由學院填寫)	※ 以本級第4項(3)、(4)申請者，「傑出績效說明表」請先至產學處確認核章後，再送回各系所審查。

# 論文方面

以前一年度(曆制年)發表論文點數為申請門檻者，需至近五年內傑出績效表填寫(同理適用產學、技轉等同項的格子)

國立臺北科技大學獎勵特殊優秀研究人才近五年內之傑出績效說明表

學術論著		年度					小計 (A)
		108	109	110	111	112	
Scopus 或 WOS 資料庫	篇數						
	點數 (請參照表 B)						
	說明：採計 Scopus 論文者，請檢附-表 B「傑出論文績效說明表」						
TSSCI/THCI (限設計及 人社學院)	篇數						
	點數 (2點/篇)						
人文、設計、藝術或 社會之學術專書	冊數						
	點數 (6點/冊)						
人文、設計、藝術或 社會之學術專書單篇 (章)	篇數						
	點數 (2點/篇)						

A

申請項目(請勾選並檢附-表 A「近五年內之傑出績效說明表」) ※本次補助起始日:113.8.1

第一級 至 第四級	申請級數：第_____級 第_____項 適用標準：_____ ※ 請依「附表一：本校獎勵特殊優秀研究人才支給標準表」之等級適用標準自行填寫
第五級	<input type="checkbox"/> 1.國際知名之國家院士 <input type="checkbox"/> 2.曾(現)任國際著名大學之講座 <input type="checkbox"/> 3.前一年度以本校名義獲國科會傑出研究獎且累積獲獎次數達二次以上 <input type="checkbox"/> 4.當學年度獲聘為本校終身講座教授者 <input type="checkbox"/> 5.前一年度以本校名義發表之論文點數達250點以上(不含研討會論文)
第六級	<input type="checkbox"/> 1.前一年度以本校名義獲國科會傑出研究獎 <input type="checkbox"/> 2.當學年度獲聘為本校專任講座教授者 <input type="checkbox"/> 3.前一年度以本校名義發表之論文點數達140點以上(不含研討會論文)
第七級	<input type="checkbox"/> 1.當學年度獲聘為本校終身特聘教授者 <input type="checkbox"/> 2.前一年度獲本校傑出研究獎或傑出產學合作獎且累積次數達二次以上者 <input type="checkbox"/> 3.前一年度以本校名義發表之論文點數達80點以上(不含研討會論文) <input type="checkbox"/> 4.前一年度以本校名義所獲得之產學合作計畫點數達240點以上且管理費納入校務基金超過150萬元 <input type="checkbox"/> 5.前一年度以本校名義所獲得之實收技術移轉金點數達175點以上且管理費納入校務基金超過50萬元 ※ 以本級第4、5項申請者，「傑出績效說明表」請先至產學處確認核章後，再送回各系所審查。
第八級 及 第九級	<input type="checkbox"/> 1.當學年度獲聘為本校特聘教授者 <input type="checkbox"/> 2.前一年度獲本校傑出研究獎者 <input type="checkbox"/> 3.前一年度獲本校傑出產學合作獎者 <input type="checkbox"/> 4.基本門檻：近五年內曾主持國科會各型計畫，並滿足下列條件之一者 <ul style="list-style-type: none"> <li><input type="checkbox"/> (1)近五年以本校名義發表之重要學術論著績效點數12點。 設計學院及人社學院教師得採計 TSSCI/THCI 期刊論文；人文、設計、藝術或社會科學領域教師得以學術專書著作或專章申請。</li> <li><input type="checkbox"/> (2)以本校名義主持國科會各類型計畫，五年內之總金額，以五年內之計畫總金額為通過標準，其標準由各學院依相關程序訂定之。</li> <li><input type="checkbox"/> (3)近五年以本校名義所獲得之產學合作計畫累計總金額超過1000萬元(績效點數200點)且管理費納入校務基金超過150萬元者。</li> <li><input type="checkbox"/> (4)近五年以本校名義所獲之實收技術移轉金累計總金額超過250萬元(績效點數125點)且管理費納入校務基金超過50萬元者。</li> </ul>
申請第4項 (1)(2)院推薦 排序	
(由學院填寫)	※ 以本級第4項(3)、(4)申請者，「傑出績效說明表」請先至產學處確認核章後，再送回各系所審查。

## 產學與技轉方面

以產學合作計畫或技術移轉金申請者，請先由產學處確認填報數據正確(以產學合作計畫為例)

第七級-4. 以前一年度申請者：

產學合作計畫 (不包含以學校名義開辦訓練課程招生收入)		年度					小計
		108	109	110	111	112	
近五年以本校名義所獲得之產學合作計畫，其實際納入本校校務基金之統計表	件數						
	管理費(萬元)						
	計畫金額(萬元)						
	點數(2點/10萬元)						
	產學處(簽章)	承辦人	單位主管				
※以第七級第4項、第八級及第九級第4項(3)申請者，請產學處填寫確認計畫及管理費總金額後核章							

第八、九級-4(3). 近五年內(加總)以本校名義申請者：

技術移轉金 (不包含國科會先期技術移轉授權金)		年度					小計
		108	109	110	111	112	
近五年以本校名義所獲之實收技術移轉金統計表	件數						
	管理費(萬元)						
	技轉金額(萬元)						
	點數(5點/10萬元)						
	產學處(簽章)	承辦人	單位主管				
※以第七級第5項、第八級及第九級第4項(4)申請者，請產學處填寫確認技轉金及管理費總金額後核章							

申請項目(請勾選並檢附-表 A「近五年內之傑出績效說明表」) ※本次補助起始日:113.8.1

第一級至第四級

申請級數：第\_\_\_\_\_級 第\_\_\_\_\_項

適用標準：\_\_\_\_\_

※ 請依「附表一：本校獎勵特殊優秀研究人才支給標準表」之等級適用標準自行填寫

第五級

- 1.國際知名之國家院士
- 2.曾(現)任國際著名大學之講座
- 3.前一年度以本校名義獲國科會傑出研究獎且累積獲獎次數達二次以上
- 4.當學年度獲聘為本校終身講座教授者
- 5.前一年度以本校名義發表之論文點數達250點以上(不含研討會論文)

第六級

- 1.前一年度以本校名義獲國科會傑出研究獎
- 2.當學年度獲聘為本校專任講座教授者
- 3.前一年度以本校名義發表之論文點數達140點以上(不含研討會論文)

第七級

- 1.當學年度獲聘為本校終身特聘教授者
  - 2.前一年度獲本校傑出研究獎或傑出產學合作獎且累積次數達二次以上者
  - 3.前一年度以本校名義發表之論文點數達80點以上(不含研討會論文)
  - 4.前一年度以本校名義所獲得之產學合作計畫點數達240點以上且管理費納入校務基金超過150萬元
  - 5.前一年度以本校名義所獲得之實收技術移轉金點數達175點以上且管理費納入校務基金超過50萬元
- ※ 以本級第4、5項申請者，「傑出績效說明表」請先至產學處確認核章後，再送回各系所審查。

第八級及第九級

- 1.當學年度獲聘為本校特聘教授者
- 2.前一年度獲本校傑出研究獎者
- 3.前一年度獲本校傑出產學合作獎者
- 4.基本門檻：近五年內曾主持國科會各型計畫，並滿足下列條件之一者
  - (1)近五年以本校名義發表之重要學術論著績效點數12點。設計學院及人社學院教師得採計 TSSCI/THCI 期刊論文；人文、設計、藝術或社會科學領域教師得以學術專書著作或專章申請。
  - (2)以本校名義主持國科會各類型計畫，五年內之總金額，以五年內之計畫總金額為通過標準，其標準由各學院依相關程序訂定之。
  - (3)近五年以本校名義所獲得之產學合作計畫累計總金額超過1000萬元(績效點數200點)且管理費納入校務基金超過150萬元者。
  - (4)近五年以本校名義所獲之實收技術移轉金累計總金額超過250萬元(績效點數125點)且管理費納入校務基金超過50萬元者。

申請第4項(1)(2)院推薦排序

(由學院填寫)

※ 以本級第4項(3)、(4)申請者，「傑出績效說明表」請先至產學處確認核章後，再送回各系所審查。

## 國科會計畫方面

- 需提供佐證資料(教師評鑑系統列印畫面)
- 通過標準由各學院訂定

國科會計畫 (不包含國科會產學合作計畫)		年度					小計
		108	109	110	111	112	
近五年以本校名義主持國科會各類型計畫統計表  (請系所與學院依教評系統資料確認)	件數						
	計畫金額 (萬元)						
	點數 (5點/10萬元)						

申請項目(請勾選並檢附-表 A「近五年內之傑出績效說明表」) ※本次補助起始日:113.8.1	
第一級 至 第四級	申請級數：第_____級 第_____項 適用標準：_____ ※ 請依「附表一：本校獎勵特殊優秀研究人才支給標準表」之等級適用標準自行填寫
第五級	<input type="checkbox"/> 1.國際知名之國家院士 <input type="checkbox"/> 2.曾(現)任國際著名大學之講座 <input type="checkbox"/> 3.前一年度以本校名義獲國科會傑出研究獎且累積獲獎次數達二次以上 <input type="checkbox"/> 4.當學年度獲聘為本校終身講座教授者 <input type="checkbox"/> 5.前一年度以本校名義發表之論文點數達250點以上(不含研討會論文)
第六級	<input type="checkbox"/> 1.前一年度以本校名義獲國科會傑出研究獎 <input type="checkbox"/> 2.當學年度獲聘為本校專任講座教授者 <input type="checkbox"/> 3.前一年度以本校名義發表之論文點數達140點以上(不含研討會論文)
第七級	<input type="checkbox"/> 1.當學年度獲聘為本校終身特聘教授者 <input type="checkbox"/> 2.前一年度獲本校傑出研究獎或傑出產學合作獎且累積次數達二次以上者 <input type="checkbox"/> 3.前一年度以本校名義發表之論文點數達80點以上(不含研討會論文) <input type="checkbox"/> 4.前一年度以本校名義所獲得之產學合作計畫點數達240點以上且管理費納入校務基金超過150萬元 <input type="checkbox"/> 5.前一年度以本校名義所獲得之實收技術移轉金點數達175點以上且管理費納入校務基金超過50萬元 ※ 以本級第4、5項申請者，「傑出績效說明表」請先至產學處確認核章後，再送回各系所審查。
第八級 及 第九級	<input type="checkbox"/> 1.當學年度獲聘為本校特聘教授者 <input type="checkbox"/> 2.前一年度獲本校傑出研究獎者 <input type="checkbox"/> 3.前一年度獲本校傑出產學合作獎者 <input type="checkbox"/> 4.基本門檻：近五年內曾主持國科會各型計畫，並滿足下列條件之一者 <ul style="list-style-type: none"> <li><input type="checkbox"/> (1)近五年以本校名義發表之重要學術論著績效點數12點。 設計學院及人社學院教師得採計 TSSCI/THCI 期刊論文；人文、設計、藝術或社會科學領域教師得以學術專書著作或專章申請。</li> <li><input type="checkbox"/> (2)以本校名義主持國科會各類型計畫，五年內之總金額，以五年內之計畫總金額為通過標準，其標準由各學院依相關程序訂定之。</li> <li><input type="checkbox"/> (3)近五年以本校名義所獲得之產學合作計畫累計總金額超過1000萬元(績效點數200點)且管理費納入校務基金超過150萬元者。</li> <li><input type="checkbox"/> (4)近五年以本校名義所獲之實收技術移轉金累計總金額超過250萬元(績效點數125點)且管理費納入校務基金超過50萬元者。</li> </ul>
申請第4項 (1)(2)院推薦 排序	<input type="checkbox"/> (1)近五年以本校名義所獲得之產學合作計畫累計總金額超過1000萬元(績效點數200點)且管理費納入校務基金超過150萬元者。 <input type="checkbox"/> (2)近五年以本校名義所獲之實收技術移轉金累計總金額超過250萬元(績效點數125點)且管理費納入校務基金超過50萬元者。 ※ 以本級第4項(3)、(4)申請者，「傑出績效說明表」請先至產學處確認核章後，再送回各系所審查。
(由學院填寫)	※ 以本級第4項(3)、(4)申請者，「傑出績效說明表」請先至產學處確認核章後，再送回各系所審查。

# 【共通】研究彈薪申請表論文面向-A1

國立臺北科技大學獎勵特殊優秀研究人才近五年內之傑出績效說明表

A

學術論著		年度					小計 (A)
		108	109	110	111	112	
Scopus 或 WOS 資料庫	篇數						
	點數 (請參照表 B)						
	說明：採計 Scopus 論文者，請檢附-表 B「傑出論文績效說明表」						
TSSCI/THCI (限設計及 人社學院)	篇數						
	點數 (2點/篇)						
人文、設計、藝術或 社會之學術專書	冊數						
	點數 (6點/冊)						
人文、設計、藝術或 社會之學術專書單篇 (章)	篇數						
	點數 (2點/篇)						

填寫以當年度往前推5年的資料並加總  
推薦以scopus的查詢為主  
須提供scopus頁面+論文封面佐證

- 專書與單篇須為初版，再版、二三刷皆不可再計入
- 專書單篇以篇數計，N篇得N\*2點

# 【共通】研究彈薪申請表論文面向-A2

\*以112年申請資料為例

申請人於 SciVal 資料庫中 **近五年** FWCI 值及 h-5 指數，若為本校近五年 FWCI 值及 h-5 指數之倍數，擇最優一項加計點數，對應表如下：

FWCI 之倍數	1.1-1.3(不含)	1.3-1.5(不含)	1.5-1.8(不含)	1.8-2.2(不含)	2.2以上
h-5指數之倍數	0.10-0.15(不含)	0.15-0.25(不含)	0.25-0.40(不含)	0.40-0.55(不含)	0.55以上
加計點數	6	8	10	13	15

申請人近五年 FWCI 值： 為本校近五年 FWCI 值 **1.02** 之 倍。

申請人 h-5 指數： 為本校 h-5 指數 **64** 之 此值以本處查詢結果為準

上述兩者擇最優一項，加計點數： 點(B)。

總計點數  
(A)+ (B)

說明：1. 近五年以本校名義發表之學術論著（此段期間曾生產或請育嬰假者得以延長，其延長期限依實際請假時間為依據，並檢附相關證明文件）始得採計。

2. 論文之期刊排名以出版年度為準，若無該出版年資料，則以前一年度為準。

3. 每篇論文僅能單一作者提出申請，若有2位或以上本校教師為共同作者，請檢附其他教師同意書。

Chen, Shenming ☆

TWN National Taipei University of Technology ... Show all affiliations | View in Scopus | Is this you?

2018 to 2022

All subject areas

ASJC

需查詢近五年之資料(EX:112年度申請，此區間應為2018 to 2022)

Summary Topics Collaboration Published Viewed Cited Patent Impact

Overall research performance

592

Scholarly Output ⓘ

20.9% All Open Access

View list of publications

1.60

Field-Weighted Citation Impact ⓘ

Yearly breakdown

12,162

Citation Count ⓘ

20.5

Citations per Publication ⓘ

83

h-index ⓘ

41

h5-index ⓘ

以陳生明教授為例：

申請人近五年FWCI值：1.6為本校近五年FWCI值 1.02 之 1.56倍 (10點)

申請人h-5指數：41 為本校h-5指數 64 之 0.64倍 (15點)

上述兩者擇最優一項，加計點數： 15點

\*操作方式詳後

# 【共通】研究彈薪申請表論文面向-A3

## 查詢方式:

登入Scopus資料庫，輸入老師名字後，點選【匯出至SciVal】，即跑出資料頁面



**Chen, Shenming -m**

National Taipei University of Technology, Taipei, Taiwan | 57449811900 | 連接到 ORCID | 這是您嗎? 連線到 Mendeley 帳戶 展開

38,495 引文 by 24,089 文獻 | 1,249 文獻 | 83 h-索引 查閱 h-圖表 | View all metrics >

設定新通報 | 儲存至清單 | 編輯作者檔案 | ... 更多

文獻與引用趨勢 | 147 | 6,291

可能比對到的作者

匯出至 SciVal



SciVal | Overview | Benchmarking | Collaboration | Trends | Reporting | My SciVal | Scopus >

**Chen, Shenming** ☆

TWN National Taipei University of Technology ... Show all affiliations | View in Scopus > | Is this you?

2018 to 2022 | All subject areas | ASJC | [Toggle]

Summary | Topics | Collaboration | Published | Viewed | Cited | Patent Impact

Overall research performance


592 Scholarly Output ⓘ 20.9% All Open Access View list of publications	1.60 Field-Weighted Citation Impact ⓘ Yearly breakdown	12,162 Citation Count ⓘ
20.5 Citations per Publication ⓘ	83 h-index ⓘ 16	41 h5-index ⓘ



# 【共通】研究彈薪申請表論文面向-A4

更改查詢區間方式:

Step3:點選右上角【Benchmarking】頁籤後，點選【Table】



Overview **Benchmarking** Collaboration Trends Reporting My SciVal Scopus 1


## Benchmarking

2018 to 2022  All subject areas  ASJC

**All Metrics** Rankings Metrics

Benchmark one metric over time | Benchmark multiple metrics   Heatmap

Step2:點選箭頭後，即跳出視窗，輸入近五年之年度後，點選【Apply】



## Benchmarking

2018 to 2022  All subject areas  ASJC

## Benchmarking

2018 to 2022  All subject areas

From:  to

1996  2018 2022 >2023

# 【共通】研究彈薪申請表論文面向-A5

更改查詢區間方式:

Step3:點選右上角【+1】頁籤後，下拉選單點選【Published】後，即出現【h-indices】，選擇【h5-index】後，點選【Choose metric】。

Step4:同前一步驟，下拉選單點選【Cited】後，即出現【Field-Weighted Citation Impact】，點選【Choose metric】。

Remove all Scholarly Output Field-Weighted Citation Impact +1

593 1.57

h-indices

A measure of both the productivity and citation impact of an entity, based on the number of publications as well as the number of citations they have received. h5-index is h-index for the past 5 years. g-index emphasizes the most highly cited publications. m-index is h-index per year.

Learn more about this metric

Select metric:

h-index  h5-index  g-index  m-index

Choose metric >

Collaboration Published Scholarly Output Subject Area Count Scopus Source Title Count h-indices Viewed Cited Patent Impact Media Impact Topic Indicators Awarded Grants

Field-Weighted Citation Impact h5-index +1

593 1.57 41

Field-Weighted Citation Impact

The ratio of citations received relative to the expected world average for the subject field, publication type and publication year. The world average FWCI is 1.00.

Note: for entities with a small scholarly output, please beware of highly cited publications which may skew the FWCI.

Learn more about this metric

Include self-citations

Choose metric >

Collaboration Published Viewed Cited Citation Count Field-Weighted Citation Impact Outputs in Top Citation Percentiles Publications in Journal Quartiles Publications in Top Journal Percentiles Citations per Publication Cited Publications

Entity ↑ Remove all Scholarly Output Field-Weighted Citation Impact h5-index +1

Chen, Shenming -m 593 1.57 41

# 【共通】研究彈薪申請表論文面向-B1

國立臺北科技大學傑出論文績效說明表

B

申請人姓名(中/英文):

系所/職稱:

員工編號:

以莊賀喬教授之論文為例:  
(接續下頁)

每篇論文僅能有一位作者提出申請，若有2位以上本校教師為**共同作者**，請檢附**其他教師同意書**

Journal Papers 請依序填寫：姓名、著作名稱、期刊名稱、卷數、頁數、發表年份(SCI/SSCI, Impact Factor, Scopus CiteScore Rank, 領域別) 並以*註記該篇所有之通訊作者，檢附每篇論文首頁與以 Scopus 資料庫為主之證明文件。	期刊排名 R (W1)	作者排序 (W2)	共同作者數 (W3)	額外加權 (W4)	國際合著學術機構國家數 (W5)	點數 (=W1×W2×W3×W4×W5)
<b>範例</b> AAA, <u>BBB*</u> , CCC, An entry-exit path planner for an autonomous tractor in a paddy field, Computers and Electronics in Agriculture, Vol.191, Dec, 2021. (SCI, IF=6.757; CiteScore Rank: 1/94=0.0106=1.06%, Horticulture)	<input type="checkbox"/> Nature、Science 及 Cell (150點) <input type="checkbox"/> R ≤ 1% (40點) <input checked="" type="checkbox"/> 1% < R ≤ 5% (25點) <input type="checkbox"/> 5% < R ≤ 10% (15點) <input type="checkbox"/> 10% < R ≤ 25% (10點) <input type="checkbox"/> 25 < R ≤ 40% (5點) <input type="checkbox"/> R > 40% (2點)	<input type="checkbox"/> 第一作者(x1) <input checked="" type="checkbox"/> 通訊作者(x1) <input type="checkbox"/> 第二作者(x0.8) <input type="checkbox"/> 第三作者(x0.6) <input type="checkbox"/> 第四作者(x0.4) <input type="checkbox"/> 第五作者以上(x0.2)	<input type="checkbox"/> 無(x1) <input checked="" type="checkbox"/> 1位通訊作者(x1) <input type="checkbox"/> 2位(含)以上通訊作者(x0.8) <input type="checkbox"/> 有多位作者 Equal Contribution (x0.8)	<input checked="" type="checkbox"/> 無 (x1) <input type="checkbox"/> 企業 (x1.1) <input type="checkbox"/> SDG (x1.1) <input type="checkbox"/> SSCI (x1.5) <input type="checkbox"/> 企業、SDG (x1.2) <input type="checkbox"/> 企業、SSCI (x1.6) <input type="checkbox"/> SDG、SSCI (x1.6) <input type="checkbox"/> 企業、SDG、SSCI (x1.8)	<input checked="" type="checkbox"/> 無 (x1) <input type="checkbox"/> 1-2個國家 (x1.1) <input type="checkbox"/> 3個國家以上 (x1.2)	25×1×1×1×1 = 25
Shobana Sebastin Mary Manickaraj, Sabarison Pandiyarajan, Ai-Ho Liao, Atchaya Ramachandran, Sheng-Tung Huang, Priyadharshini Natarajan, <u>Ho-Chiao Chuang*</u> , “Sansevieria trifasciata biomass-derived activated carbon by supercritical-CO <sub>2</sub> route: electrochemical detection towards carcinogenic organic pollutant and energy storage application” Electrochimica Acta, Vol.424, pp 140672, August 2022. (SCI, Impact Factor=7.3; CiteScore Rank: 19/280=6.78%, General Chemical Engineering)	<input type="checkbox"/> Nature、Science 及 Cell (150點) <input type="checkbox"/> R ≤ 1% (40點) <input type="checkbox"/> 1% < R ≤ 5% (25點) <input checked="" type="checkbox"/> 5% < R ≤ 10% (15點) <input type="checkbox"/> 10% < R ≤ 25% (10點) <input type="checkbox"/> 25 < R ≤ 40% (5點) <input type="checkbox"/> R > 40% (2點)	<input type="checkbox"/> 第一作者(x1) <input type="checkbox"/> 通訊作者(x1) <input type="checkbox"/> 第二作者(x0.8) <input type="checkbox"/> 第三作者(x0.6) <input type="checkbox"/> 第四作者(x0.4) <input type="checkbox"/> 第五作者以上(x0.2)	<input type="checkbox"/> 無(x1) <input type="checkbox"/> 1位通訊作者(x1) <input type="checkbox"/> 2位(含)以上通訊作者(x0.8) <input type="checkbox"/> 有多位作者 Equal Contribution (x0.8)	<input type="checkbox"/> 無(x1) <input type="checkbox"/> 企業 (x1.1) <input type="checkbox"/> SDG (x1.1) <input type="checkbox"/> SSCI (x1.5) <input type="checkbox"/> 企業、SDG (x1.2) <input type="checkbox"/> 企業、SSCI (x1.6) <input type="checkbox"/> SDG、SSCI (x1.6) <input type="checkbox"/> 企業、SDG、SSCI (x1.8)	<input type="checkbox"/> 無 (x1) <input type="checkbox"/> 1-2個國家 (x1.1) <input type="checkbox"/> 3個國家以上 (x1.2)	15 * 1 * 1 * 1.1 * 1.1 = 18.15

續下頁

# 【共通】研究彈薪申請表論文面向-B2

查詢方式: (以莊賀喬教授之論文為例)

Step1: 登入Scopus資料庫

(<https://www.scopus.com/search/form.uri?display=authorLookup#basic>)，輸入老師名字後，點選【搜尋】

開始探索  
探索最可靠、最相關、最即時的研究，一站式處理。

文獻 作者 搜尋研究人員 (Researcher Discovery) 機構 Scopus AI *Alpha* 搜尋提示

Search authors using:  作者姓名  ORCID  關鍵字 新增

輸入姓氏 \*  
Chuang

輸入名字  
Ho-Chiao

+ 新增機構

搜尋

Step2: 確認所屬機構為本校後，點選【老師名字】

僅顯示完全相符

優化搜尋結果

限制範圍 排除

機構

- National Taipei University of Technology (1) >
- National Tsing Hua University (1) >
- University of Colorado Boulder (1) >

城市

排序方式: 文獻數量 (高至低)

全部 顯示文獻 Citation overview 請求合併作者

作者	文獻	h-index	機構	城市	國家/地區
<input type="checkbox"/> 1 Chuang, Hochiao Chiao Rick Chuang, Ho Chiao Chuang, Chiao Ho Chuang, H. C.	103	14	National Taipei University of Technology	Taipei	Taiwan

查看最近的文獻標題

每頁顯示: 20 個搜尋結果 / 每頁 1 頁首

20

續下頁

# 【共通】研究彈薪申請表論文面向-B3

Step3:點入之後滑至最下方，將顯示調至【200結果】

### Chuang, Hochiao Chiao Rick

National Taipei University of Technology, Taipei, Taiwan © 54083059900 連接到 ORCID 展開

627 103 14  
引用 by 414 文獻 文獻 h-索引 查閱 h-圖表 View all metrics >

設定新通知 編輯作者檔案 更多

#### 文獻與引用趨勢

貢獻度最多的主題 2018–2022

- Supercritical CO<sub>2</sub>; Supercritical Carbon Dioxide; Difluoromethane 13 篇文獻
- Four-Dimensional Computed Tomography; Fiducial Markers; Cancer 8 篇文獻
- Air Conditioning; Evaporators; Cold Chain 4 篇文獻

查看所有主題

103 文獻 作者指標 Cited by 414 文獻 1 預印本 162 共同作者 20 主題 0 獎勵補助款

Article

### 3D-flower-like porous neodymium molybdate nanostructure for trace level detection of organophosphorus pesticide in food samples

Ganesan, M., Keerthika Devi, R., Liao, A.-H., ...Gopalakrishnan, G., Chuang, H.-C.

*Food Chemistry*, 2022, 396, 133722

8 8  
引用次數

查看摘要 查看摘要 查看摘要 查看摘要 查看摘要 查看摘要 查看摘要 查看摘要 查看摘要 查看摘要

NTUT Full-Text Full Text 相關文獻

< 上一頁 1 2 3 4 5 ... 11 下一頁 >

顯示 10 結果

- 10 結果
- 20 結果
- 50 結果
- 100 結果
- 200 結果

opus 語言 客戶服務

# 【共通】研究彈薪申請表論文面向-B4

Step4:利用CTRL+F去快速搜尋本篇論文

The screenshot shows a search results page for the author 'Chuang, Hochiao Chiao Rick'. The browser's address bar is highlighted with a red box, showing the search term 'Sansevieria trifasciata bioma' and page '1/1'. The search results are listed in a table-like format with columns for article title, citation count, and actions. The third article is highlighted with a red box.

Article	引用次數
Combining the wavelet transform with a phase-lead compensator to a respiratory motion compensation system with an ultrasound tracking technique in radiation therapy Kuo, C.-C., Guo, M.-L., Liao, A.-H., ...Ting, L.-L., Chuang, H.-C. <i>Biomedical Signal Processing and Control</i> , 2022, 78, 103892 查看摘要 <input type="button" value="NTUT Full-Text"/> <input type="button" value="Full Text"/> 相關文獻	0
Malic acid pathway of constructing high-performance Ni anticorrosive coatings using supercritical-CO <sub>2</sub> electrodeposition Manickaraj, S.S.M., Pandiyarajan, S., Liao, A.-H., ...Lee, K.-Y., Chuang, H.-C. <i>Materials Science In Semiconductor Processing</i> , 2022, 148, 106780 查看摘要 <input type="button" value="NTUT Full-Text"/> <input type="button" value="Full Text"/> 相關文獻	5
<b>Sansevieria trifasciata biomass-derived activated carbon by supercritical-CO<sub>2</sub> route: Electrochemical detection towards carcinogenic organic pollutant and energy storage application</b> Manickaraj, S.S.M., Pandiyarajan, S., Liao, A.-H., ...Natarajan, P., Chuang, H.-C. <i>Electrochimica Acta</i> , 2022, 424, 140672 查看摘要 <input type="button" value="NTUT Full-Text"/> <input type="button" value="Full Text"/> 相關文獻	6

# 【共通】研究彈薪申請表論文面向-B5

Step5:點選反橘色之論文題目，即帶入論文資料畫面，要確認論文發布時間在本次申請之規定時間內

*Electrochimica Acta* · 卷 424 · **20 August 2022** · 論文號碼 140672

文獻類型  
論文  
來源出版物種類  
期刊  
ISSN :  
00134686  
DOI  
10.1016/j.electacta.2022.140672  
展開

## Sansevieria trifasciata biomass-derived activated carbon by supercritical-CO<sub>2</sub> route: Electrochemical detection towards carcinogenic organic pollutant and energy storage application

Manickaraj, Shobana Sebastin Mary<sup>a, b</sup>; Pandiyarajan, Sabarison<sup>a, b</sup>;  
Liao, Ai-Ho<sup>c, d</sup>; Ramachandran, Atchaya<sup>e</sup>; Huang, Sheng-Tung<sup>a</sup>;  
Natarajan, Priyadharshini<sup>f</sup>; Chuang, Ho-Chiao<sup>b</sup> ✉  
[將全部儲存到作者清單](#)

<sup>a</sup> Department of Chemical Engineering and Biotechnology, National Taipei University of Technology, Taipei, 106344, Taiwan  
<sup>b</sup> Department of Mechanical Engineering, National Taipei University of Technology, Taipei, 106344, Taiwan  
<sup>c</sup> Graduate Institute of Biomedical Engineering, National Taiwan University of Science and Technology, Taipei, 106335, Taiwan  
<sup>d</sup> Department of Biomedical Engineering, National Defense Medical Center, Taipei, 114201, Taiwan  
[顯示其他的機構](#)

6 78th percentile  
在 Scopus 中的引用次數 : in Scopus

1.26  
領域權重引用影響指數 (FWCI)

14  
瀏覽次數

[查看所有計量](#)

[查閱 PDF](#) [全文選項](#) [匯出](#)

摘要  
作者關鍵字  
Reaxys 化學資料庫資訊

摘要  
Activated carbon (AC) has been widely used for electrochemical applications, such as electrochemical sensors, energy storage applications, etc., due to its fine porous structure, volumetric capacitance, and chemical stability. Supercritical-CO<sub>2</sub> (SC-CO<sub>2</sub>) has a fascinating advantage in material science due to its microbubble cavitation, high diffusivity, and high

# 【共通】研究彈薪申請表論文面向-B6

**B**

查詢W1~W5之方式

## 國立臺北科技大學傑出論文績效說明表

申請人姓名(中/英文)：

系所/職稱：

員工編號：

Journal Papers 請依序填寫：姓名、著作名稱、期刊名稱、卷數、頁數、發表年份(SCI/SSCI,Impact Factor,Scopus CiteScore Rank,領域別) 並以*註記該篇所有之通訊作者，檢附每篇論文首頁與以 Scopus 資料庫為主之證明文件。		期刊排名 R (W1)	作者排序 (W2)	共同作者數 (W3)	額外加權 (W4)	國際合著學術機構國家數 (W5)	點數 (=W1×W2×W3×W4×W5)
1	<input type="checkbox"/> Nature、Science 及 Cell (150點) <input type="checkbox"/> $R \leq 1\%$ (40點) <input type="checkbox"/> $1\% < R \leq 5\%$ (25點) <input type="checkbox"/> $5\% < R \leq 10\%$ (15點) <input type="checkbox"/> $10\% < R \leq 25\%$ (10點) <input type="checkbox"/> $25 < R \leq 40\%$ (5點) <input type="checkbox"/> $R > 40\%$ (2點)	<input type="checkbox"/> 第一作者(x1) <input type="checkbox"/> 通訊作者(x1) <input type="checkbox"/> 第二作者(x0.8) <input type="checkbox"/> 第三作者(x0.6) <input type="checkbox"/> 第四作者(x0.4) <input type="checkbox"/> 第五作者以上(x0.2)	<input type="checkbox"/> 無(x1) <input type="checkbox"/> 1位通訊作者(x1) <input type="checkbox"/> 2位(含)以上通訊作者(x0.8) <input type="checkbox"/> 有多位作者 Equal Contribution (x0.8)	<input type="checkbox"/> 無(x1) <input type="checkbox"/> 企業 (x1.1) <input type="checkbox"/> SDG (x1.1) <input type="checkbox"/> SSCI (x1.5) <input type="checkbox"/> 企業、SDG (x1.2) <input type="checkbox"/> 企業、SSCI (x1.6) <input type="checkbox"/> SDG、SSCI (x1.6) <input type="checkbox"/> 企業、SDG、SSCI (x1.8)	<input type="checkbox"/> 無 (x1) <input type="checkbox"/> 1-2個國家 (x1.1) <input type="checkbox"/> 3個國家以上 (x1.2)	24	



# 【共通】研究彈薪申請表論文面向-B7

查詢W1方式-以Scopus查詢

Step1:點選期刊名稱後，視窗右邊即顯示出來源出版物詳情預覽欄位，點選【瀏覽完整的來源出版物詳情】

**Electrochimica Acta** · 卷 424 · 20 August 2022 · 論文號碼 140672

## Sansevieria trifasciata biomass-derived activated carbon by supercritical-CO<sub>2</sub> route: Electrochemical detection towards carcinogenic organic pollutant and energy storage application

Manickaraj, Shobana Sebastin Mary<sup>a, b</sup>; Pandiyarajan, Sabarison<sup>a, b</sup>; Liao, Ai-Ho<sup>c, d</sup>; Ramachandran, Atchaya<sup>e</sup>; Huang, Sheng-Tung<sup>a</sup>; Natarajan, Priyadharshini<sup>f</sup>; Chuang, Ho-Chiao<sup>b</sup> ✉

[將全部儲存到作者清單](#)

<sup>a</sup> Department of Chemical Engineering and Biotechnology, National Taipei University of Technology, Taipei, 106344, Taiwan  
<sup>b</sup> Department of Mechanical Engineering, National Taipei University of Technology, Taipei, 106344, Taiwan  
<sup>c</sup> Graduate Institute of Biomedical Engineering, National Taiwan University of Science and Technology, Taipei, 106335, Taiwan  
<sup>d</sup> Department of Biomedical Engineering, National Defense Medical Center, Taipei, 114201, Taiwan

[顯示其他的機構](#) ▾

6 78th percentile  
在 Scopus 中的引用次數: in Scopus

1.26  
領域權重引用影響指數 (FWCI)

14  
瀏覽次數

[查看所有計量](#) >

[查閱 PDF](#) [全文選項](#) ▾ [匯出](#) ▾

來源出版物詳情預覽

### Electrochimica Acta

出版: Elsevier  
來源出版物種類: 期刊

[瀏覽完整的來源出版物詳情](#)

---

#### 計量

12.8 CiteScore 2022 ⓘ	1.264 SJR 2022 ⓘ	1.047 SNIP 2022 ⓘ
--------------------------	---------------------	----------------------

#### CiteScore 排名 ⓘ

ASJC 類別	四分位數	百分位數	排名
General Chemical Engineering	Q1	93rd	19 / 272
Electrochemistry	Q1	87th	7 / 54

25

# 【共通】研究彈薪申請表論文面向-B8

查詢W1方式-以Scopus查詢

Step2:選擇論文發表時的年份(如2022年發表，則應選擇2022年之CiteScore)

來源出版物詳情 回饋 > 比較來源出版物 >

Electrochimica Acta  
Scopus 涵蓋年度: 從 1959 至今  
圖書館訂閱: 從 January 1995  
發表者: Elsevier  
國際標準期刊號: 0013-4686  
學科類別: Chemical Engineering: General Chemical Engineering Chemistry: Electrochemistry  
來源出版物種類 期刊

[查閱所有文獻 >](#) [設定文獻通知](#) [儲存到來源出版物清單](#) [Source Homepage](#) [NTUT Full-Text](#) [Journal Finder](#) [更多 >](#)

CiteScore CiteScore 趨勢 Scopus 內容涵蓋範圍

**i** CiteScore 的改良計算方法 ×

CiteScore 2022 計算在 2019-2022發表的論文、回顧文獻、會議論文、專書論文、和數據論文等等在 2019-2022 所收到的引用總數，除以發表於2019-2022的出版物總數。 [瞭解更多 >](#)

CiteScore 2022 = 12.8

CiteScore 追蹤2023 ①

10.3 =  $\frac{\text{迄今 62,619 個引用次數}}{\text{迄今 6,087 篇文獻}}$

最後一次更新: 05 September, 2023 • 每個月更新

計算 05 May, 2023

CiteScore 排

類別

百分位數

年份	引用次數	篇文獻
2022		
2021		
2020		
2019		
2018		
2017		
2016		
2015		

# 【共通】研究彈薪申請表論文面向-B9

查詢W1方式-以Scopus查詢

Step3:取百分位數最高之排名後，將期刊排名轉換成對應點數， $19/272=6.9\%$ ，對應法規點數為15，並請檢附查詢畫面當作佐證資料



期刊排名 R (W1)
<input type="checkbox"/> Nature、Science 及 Cell (150點)
<input type="checkbox"/> $R \leq 1\%$ (40點)
<input type="checkbox"/> $1\% < R \leq 5\%$ (25點)
<input checked="" type="checkbox"/> $5\% < R \leq 10\%$ (15點)
<input type="checkbox"/> $10\% < R \leq 25\%$ (10點)
<input type="checkbox"/> $25 < R \leq 40\%$ (5點)
<input type="checkbox"/> $R > 40\%$ (2點)

# 【共通】研究彈薪申請表論文面向-B10

查詢W1方式-以WOS查詢

Step1:輸入論文題目後，點選【Search】，即帶入論文資料畫面

DOCUMENTS RESEARCHERS

Search in: Web of Science Core Collection Editions: All

DOCUMENTS CITED REFERENCES

All Fields

Example: liver disease india singh  
Sansevieria trifasciata biomass-derived activated carbon by supercritical-CO2 route: electr

+ Add row + Add date range Advanced Search

× Clear Search

0/1 Add To Marked List Export

Sort by: Relevance 1 of 1

1 Sansevieria trifasciata biomass-derived activated carbon by supercritical-CO2 route: Electrochemical detection towards carcinogenic organic pollutant and energy storage application

Manickaraj, SSM; Pandiyarajan, S; (...); Chuang, HC

Aug 20 2022 | Jun 2022 (Early Access) | ELECTROCHIMICA ACTA 424

Activated carbon (AC) has been widely used for electrochemical applications, such as electrochemical sensors, energy storage applications, etc., due to its fine porous structure, volumetric capacitance, and chemical stability. Supercritical-CO2 (SC-CO2) has a fascinating advantage in material science due to its microbubble cavitation, high diffusivity, and high permeability. In the shed of ligh... Show more

6 Citations

62 References

Full Text at Publisher

Related records

# 【共通】研究彈薪申請表論文面向-B11

查詢W1方式-以WOS查詢

Step2:點選期刊名稱後，視窗右邊即顯示出Journal information欄位，點選【Learn more】

Full text at publisher Export Add

## Sansevieria trifasciata biomass-derived activated carbon by supercritical-CO2 route: Electrochemical detection towards carcinogenic organic pollutant and energy storage application

By Manickaraj, SSM (Manickaraj, Shobana Sebastin Mary) [1], [2]; Pandiyarajan, S (Pandiyarajan, Sabarison) [1], [2]; Liao, AH (Liao, Ai-Ho) [3], [4]; Ramachandran, A (Ramachandran, Atchaya) [5]; Huang, ST (Huang, Sheng-Tung) [1]; Natarajan, P (Natarajan, Priyadharshini) [6]; Chuang, HC (Chuang, Ho-Chiao) [2]

View Web of Science ResearcherID and ORCID (provided by Clarivate)

Source **ELECTROCHIMICA ACTA**  
Volume: 424  
DOI: 10.1016/j.electacta.2022.140672

Article Number 140672

Published AUG 20 2022

Early Access JUN 2022

Indexed 2022-07-10

Document Type Article

Abstract Activated carbon (AC) has been widely used for electrochemical applications, such as electrochemical sensors, energy storage applications, etc., due to its fine porous structure, volumetric capacitance, and chemical stability. Supercritical-CO2 (SC-CO2) has a

### Journal information

ELECTROCHIMICA ACTA  
Publisher name: PERGAMON-ELSEVIER SCIENCE LTD

Journal Impact Factor™  
6.6 6  
2022 Five Year

JCR Category	Category Rank	Category Quartile
ELECTROCHEMISTRY in SCIE edition	8/30	Q2

Source: Journal Citation Reports 2022 . Learn more

Journal Citation Indicator™  
1.12 1.18  
2022 2021

JCI Category	Category Rank	Category Quartile
ELECTROCHEMISTRY in SCIE edition	7/42	Q1

The Journal Citation Indicator is a measure of the average Category Normalized Citation Impact (CNCI) of citable items (articles and reviews) published by a journal over a recent three year period. It is used to help you evaluate journals based on other metrics besides the Journal Impact Factor (JIF).

# 【共通】研究彈薪申請表論文面向-B12

查詢W1方式-以WOS查詢

Step3:選擇論文發表時的年份(如2022年發表，則應選擇2022年之JCR YEAR)

Step4:滑至中間查詢排名，取百分位數最高之排名後，將期刊排名轉換成對應點數， $7/42=16.6\%$ ，對應法規點數為10，並請檢附查詢畫面當作佐證資料

Journal Citation Reports™ Journals Categories Publishers Countries/Regions My favorites Sign In Register

Home > Journal profile Favorite Export

JCR YEAR  
2022

## ELECTROCHIMICA ACTA

ISSN  
0013-4686

EISSN  
1873-3859

JCR ABBREVIATION  
ELECTROCHIM ACTA

ISO ABBREVIATION  
Electrochim. Acta

Journal information

EDITION  
Science Citation Index Expanded (SCIE)

CATEGORY  
ELECTROCHEMISTRY - SCIE

LANGUAGES REGION 1ST ELECTRONIC JCR YEAR  
Multi-Language ENGLAND 1997

Publisher information

PUBLISHER ADDRESS PUBLICATION FREQUENCY  
PERGAMON-ELSEVIER SCIENCE LTD THE BOULEVARD, LANGFORD LANE, KIDLINGTON, OXFORD OX5 1GB, ENGLAND 28 issues/year

### Rank by Journal Impact Factor

Journals within a category are sorted in descending order by Journal Impact Factor (JIF) resulting in the Category Ranking below. A separate rank is shown for the most recent year. Data for the most recent year is presented at the top of the list, with other years shown in reverse chronological order. Learn more

EDITION  
Science Citation Index Expanded (SCIE)

CATEGORY  
ELECTROCHEMISTRY  
8/30

JCR YEAR	JIF RANK	JIF QUARTILE	JIF PERCENTILE
2022	8/30	Q2	75.0
2021	7/30	Q1	78.33
2020	8/29	Q2	74.14
2019	5/27	Q1	83.33
2018	5/26	Q1	82.69

### Rank by Journal Citation Indicator (JCI)

Journals within a category are sorted in descending order by Journal Citation Indicator (JCI) resulting in the Category Ranking below. A separate rank is shown for the most recent year. Data for the most recent year is presented at the top of the list, with other years shown in reverse chronological order. Learn more

CATEGORY  
ELECTROCHEMISTRY  
7/42

JCR YEAR	JCI RANK	JCI QUARTILE	JCI PERCENTILE
2022	7/42	Q1	84.52
2021	7/42	Q1	84.52
2020	7/39	Q1	83.33
2019	7/39	Q1	83.33
2018	6/37	Q1	85.14
2017	5/36	Q1	87.50

## 期刊排名 R (W1)

- Nature、Science 及 Cell (150點)
- $R \leq 1\%$  (40點)
- $1\% < R \leq 5\%$  (25點)
- $5\% < R \leq 10\%$  (15點)
- $10\% < R \leq 25\%$  (10點)
- $25\% < R \leq 40\%$  (5點)
- $R > 40\%$  (2點)

# 【共通】研究彈薪申請表論文面向-B13

查詢W1方式

注意事項:

1. 查詢年度應選擇**論文發表時的年份**(如2022年發表，則應選擇2022年)，倘2023年發表，因有時間差之問題，**故可先提供2022年之排名為佐證**。
2. 可自行選擇以Scopus或Wos之**查詢結果**為佐證資料。
3. 在**不四捨五入**的情況下依據其所屬區間對應**權重數值**。

# 【共通】研究彈薪申請表論文面向-B14

查詢W2方式-以Scopus查詢

依ppt第20頁方式查詢出以下畫面，點選【查閱PDF】，再點選【View PDF】即下載論文檔案

1 / 1

↓ 下載 列印 儲存至 PDF ☆ 儲存到清單 建立書目

*Electrochimica Acta* • 卷 424 • 20 August 2022 • 論文號碼 140672

文獻類型  
論文  
來源出版物種類  
期刊  
ISSN :  
00134686  
DOI  
10.1016/j.electacta.2022.140672  
展開

## Sansevieria trifasciata biomass-derived activated carbon by supercritical-CO<sub>2</sub> route: Electrochemical detection towards carcinogenic organic pollutant and energy storage application

Manickaraj, Shobana Sebastin Mary<sup>a,b</sup>; Pandiyarajan, Sabarison<sup>a,b</sup>; Liao, Ai-Ho<sup>c,d</sup>; Ramachandran, Atchaya<sup>e</sup>; Huang, Sheng-Tung<sup>a</sup>; Natarajan, Priyadharshini<sup>f</sup>; Chuang, Ho-Chiao<sup>b</sup>


<sup>a</sup> Department of Chemical Engineering and Biotechnology, National Taipei University of Technology, Taipei, 106344, Taiwan  
<sup>b</sup> Department of Mechanical Engineering, National Taipei University of Technology, Taipei, 106344, Taiwan  
<sup>c</sup> Graduate Institute of Biomedical Engineering, National Taiwan University of Science and Technology, Taipei, 106335, Taiwan  
<sup>d</sup> Department of Biomedical Engineering, National Defense Medical Center, Taipei, 114201, Taiwan

顯示其他的機構

678th percentile 在 Scopus 中的引用次數 : in Scopus	1.26 領域權重引用影響指數 (FWCI)	14 瀏覽次數	查看所有計量
---	---------------------------	------------	--------

查閱 PDF 全文選項 匯出

View PDF Download full issue

 ELSEVIER

## Electrochimica Acta

Volume 424, 20 August 2022, 140672

## Sansevieria trifasciata biomass-derived activated carbon by supercritical-CO<sub>2</sub> route: Electrochemical detection towards carcinogenic organic pollutant and energy storage application

Shobana Sebastin Mary Manickaraj<sup>a,b,1</sup>, Sabarison Pandiyarajan<sup>a,b,1</sup>, Ai-Ho Liao<sup>c,d</sup>, Atchaya Ramachandran<sup>e</sup>, Sheng-Tung Huang<sup>a</sup>, Priyadharshini Natarajan<sup>f</sup>, Ho-Chiao Chuang<sup>b</sup>

Show more

+ Add to Mendeley Share Cite

<https://doi.org/10.1016/j.electacta.2022.140672> Get rights and content



# 【共通】研究彈薪申請表論文面向-B15

查詢W2方式-以WOS查詢

依ppt第27頁方式查詢出以下畫面，點選【Full text at publisher】，再點選【View PDF】即下載論文檔案

Full text at publisher

## Sansevieria trifasciata biomass-derived activated carbon by supercritical-CO<sub>2</sub> route: Electrochemical detection towards carcinogenic organic pollutant and energy storage application

By Manickaraj, SSM (Manickaraj, Shobana Sebastin Mary) [1], [2]; Pandiyarajan, S (Pandiyarajan, Sabarison) [1], [2]; Liao, AH (Liao, Ai-Ho) [3], [4]; Ramachandran, A (Ramachandran, Atchaya) [5]; Huang, ST (Huang, Sheng-Tung) [1]; Natarajan, P (Natarajan, Priyadharshini) [6]; Chuang, HC (Chuang, Ho-Chiao) [2]

View Web of Science ResearcherID and ORCID (provided by Clarivate)

Source: ELECTROCHIMICA ACTA  
Volume: 424  
DOI: 10.1016/j.electacta.2022.140672

Article Number: 140672  
Published: AUG 20 2022  
Early Access: JUN 2022  
Indexed: 2022-07-10  
Document Type: Article

Abstract: Activated carbon (AC) has been widely used for electrochemical applications, such as electrochemical sensors, energy storage applications, etc., due to its fine porous structure, volumetric capacitance, and chemical stability. Supercritical-CO<sub>2</sub> (SC-CO<sub>2</sub>) has a fascinating advantage in material science due to its microbubble cavitation, high diffusivity, and high permeability. In the shed of light, we developed a high porous Sansevieria trifasciata biomass-derived AC by SC-CO<sub>2</sub>/SC-

View PDF

Download full issue

ELSEVIER

Electrochimica Acta  
Volume 424, 20 August 2022, 140672

## Sansevieria trifasciata biomass-derived activated carbon by supercritical-CO<sub>2</sub> route: Electrochemical detection towards carcinogenic organic pollutant and energy storage application

Shobana Sebastin Mary Manickaraj<sup>a, b, 1</sup>, Sabarison Pandiyarajan<sup>a, b, 1</sup>, Ai-Ho Liao<sup>c, d</sup>, Atchaya Ramachandran<sup>e</sup>, Sheng-Tung Huang<sup>a</sup>, Priyadharshini Natarajan<sup>f</sup>, Ho-Chiao Chuang<sup>b</sup>

Show more

+ Add to Mendeley Share Cite

<https://doi.org/10.1016/j.electacta.2022.140672> Get rights and content

# 【共通】研究彈薪申請表論文面向-B16

以陳彥霖教授之論文為例：  
本篇文章陳教授為第一作者，對應法規應x1

applied sciences MDPI

Article

## An Upper Extremity Rehabilitation System Using Efficient Vision-Based Action Identification Techniques

Yen-Lin Chen <sup>1</sup>, Chin-Hsuan Liu <sup>1</sup>, Chao-Wei Yu <sup>1</sup>, Posen Lee <sup>2,\*</sup> and Yao-Wen Kuo <sup>1</sup>

<sup>1</sup> Department Computer Science and Information Engineering, National Taipei University of Technology, Taipei 10608, Taiwan; ylchen@csie.ntut.edu.tw (Y.-L.C.); chinhsuanliu@gmail.com (C.-H.L.); david741002@gmail.com (C.-W.Y.); kent21221@gmail.com (Y.-W.K.)  
<sup>2</sup> Department Occupational Therapy, I-Shou University, Kaohsiung 82445, Taiwan  
 \* Correspondence: posenlee@isu.edu.tw; Tel.: +886-7-657-7711 (ext. 7516)

Received: 30 May 2018; Accepted: 10 July 2018; Published: 17 July 2018

**Featured Application:** This study proposes an upper extremity rehabilitation system using efficient action identification system for home based on color and depth sensor information, and can perform well under complex ambient environments.

**Abstract:** This study proposes an action identification system for home upper extremity rehabilitation. In the proposed system, we apply an RGB-depth (color-depth) sensor to capture the image sequences of the patient's upper extremity actions to identify its movements. We apply a skin color detection technique to assist with extremity identification and to build up the upper extremity skeleton points. We use the dynamic time warping algorithm to determine the rehabilitation actions. The system presented herein builds up upper extremity skeleton points rapidly. Through the upper extremity of the human skeleton and human skin color information, the upper extremity skeleton points are effectively established by the proposed system, and the rehabilitation actions of patients are identified by a dynamic time warping algorithm. Thus, the proposed system can achieve a high recognition rate of 98% for the defined rehabilitation actions for the various muscles. Moreover, the computational speed of the proposed system can reach 125 frames per second—the processing time per frame is less than 8 ms on a personal computer platform. This computational efficiency allows efficient extensibility for future developments to deal with complex ambient environments and for implementation in embedded and pervasive systems. The major contributions of the study are: (1) the proposed system is not only a physical exercise game, but also a movement training program for specific muscle groups; (2) The hardware of upper extremity rehabilitation system included a personal computer with personal computer and a depth camera. These are economic equipment, so that patients who need this system can set up one set at home; (3) patients can perform rehabilitation actions in sitting position to prevent him/her from falling down during training; (4) the accuracy rate of identifying rehabilitation action is as high as 98%, which is sufficient for distinguishing between correct and wrong action when performing specific action trainings; (5) The proposed upper extremity rehabilitation system is real-time, efficient to vision-based action identification, and low-cost hardware and software, which is affordable for most families.

**Keywords:** upper extremity identification; color and depth sensors; skeleton points; rehabilitation actions; home rehabilitation; computer vision

Yen-Lin Chen <sup>1</sup>, Chin-Hsuan Liu <sup>1</sup>, Chao-Wei Yu <sup>1</sup>, Posen Lee <sup>2,\*</sup> and Yao-Wen Kuo <sup>1</sup>

<sup>1</sup> Department Computer Science and Information Engineering, National Taipei University of Technology, Taipei 10608, Taiwan; ylchen@csie.ntut.edu.tw (Y.-L.C.); chinhsuanliu@gmail.com (C.-H.L.); david741002@gmail.com (C.-W.Y.); kent21221@gmail.com (Y.-W.K.)  
<sup>2</sup> Department Occupational Therapy, I-Shou University, Kaohsiung 82445, Taiwan  
 \* Correspondence: posenlee@isu.edu.tw; Tel.: +886-7-657-7711 (ext. 7516)

(通訊作者非陳教授)

作者排序 (W2)

第一作者(X1)

通訊作者(X1)

第二作者(X0.8)

第三作者(X0.6)

第四作者(X0.4)

第五作者以上 (X0.2)

# 【共通】研究彈薪申請表論文面向-B17

以莊賀喬教授之論文為例：  
本篇文章莊教授為通訊作者，對應法規應x1

Electrochimica Acta  
journal homepage: [www.journals.elsevier.com/electrochimica-acta](http://www.journals.elsevier.com/electrochimica-acta)

**Sansevieria trifasciata** biomass-derived activated carbon by supercritical-CO<sub>2</sub> route: Electrochemical detection towards carcinogenic organic pollutant and energy storage application

Shobana Sebastin Mary Manickaraj<sup>a,b,1</sup>, Sabarison Pandiyarajan<sup>a,b,1</sup>, Ai-Ho Liao<sup>c,d</sup>, Atchaya Ramachandran<sup>e</sup>, Sheng-Tung Huang<sup>a</sup>, Priyadharshini Natarajan<sup>f</sup>, Ho-Chiao Chuang<sup>b,\*</sup>

<sup>a</sup> Department of Chemical Engineering and Biotechnology, National Taipei University of Technology, Taipei 106344, Taiwan  
<sup>b</sup> Department of Mechanical Engineering, National Taipei University of Technology, Taipei 106344, Taiwan  
<sup>c</sup> Graduate Institute of Biomedical Engineering, National Taiwan University of Science and Technology, Taipei 106335, Taiwan  
<sup>d</sup> Department of Biomedical Engineering, National Defense Medical Center, Taipei 114201, Taiwan  
<sup>e</sup> PG & Research Department of Chemistry, Bishop Heber College, Tiruchirappalli 620017, India  
<sup>f</sup> Syngene International Pvt., Ltd. Bangalore, India

**ARTICLE INFO**

**Keywords:**  
Activated carbon  
Sansevieria trifasciata  
Supercritical-CO<sub>2</sub>  
Electrochemical sensor  
Supercapacitor

**ABSTRACT**  
Activated carbon (AC) has been widely used for electrochemical applications, such as electrochemical sensors, energy storage applications, etc., due to its fine porous structure, volumetric capacitance, and chemical stability. Supercritical-CO<sub>2</sub> (SC-CO<sub>2</sub>) has a fascinating advantage in material science due to its microbubble cavitation, high diffusivity, and high permeability. In the shed of light, we developed a high porous *Sansevieria trifasciata* biomass-derived AC by SC-CO<sub>2</sub> (SC-ST-AC). For comparison purposes, the AC was also prepared in a conventional approach (C-ST-AC). The prepared ACs were characterized through various spectroscopic and microscopic techniques to study their surface morphological character, structural analysis, and phase purity. The electrochemical performance was evaluated by two different applications: electrochemical detection and energy storage application. Based on the results, the SC-ST-AC exhibits higher porous architecture in their morphology and high phase purity with amorphous nature than C-ST-AC. In the preliminary electrochemical analysis, SC-ST-AC achieved higher performance than C-ST-AC. Thus, SC-ST-AC is applied to the real-time application and it exposed a superior limit of detection (0.005 μM L<sup>-1</sup>) and sensitivity (0.854 μA μM<sup>-1</sup> cm<sup>-2</sup>) towards MA sensing and higher specific capacitance (342.5 F/g for 2 A/g) with 92.09 % of retention at high current density. Thereby, we suggest the SC-CO<sub>2</sub> method is a promising approach to develop a highly porous carbon material with excellent electrochemical performance.

**1. Introduction**  
In recent eras, carbon-based materials including one-dimension (1D) carbon nanotubes, carbon nanofibers [1,2], two-dimension (2D) graphene [3], three-dimension (3D) graphite, activated carbon, and its derivatives [4,5] have been extensively investigated as successful commercialization materials in several sectors. Among them activated carbon (AC) is considered the most cardinal material for electrochemical application owing to its high surface area, porous architecture, and chemical stability [6-8]. The varieties of functional group moiety fascinated on the surface make it as a promising electrode material for energy storage applications [9]. Traditionally, the preparation of AC is done by the pyrolysis of fossil raw materials such as coal and petroleum coke or wood, followed by a physical or chemical activation process [10]. Due to the rapid increase of the global population and economy, the demand for energy and resources is also increasing exponentially, resulting in a lack of fossil fuels [11]. Therefore, cost-effective renewable carbon sources, the development of economic efficiency methods, and environmental safety are all issues that must be thoroughly investigated to produce advanced activated carbon that is more environmentally friendly. In this regard, biomass materials are presently recognized as the most viable candidates for preparing carbon materials

\* Corresponding author.  
E-mail address: [hchuang@mail.ntut.edu.tw](mailto:hchuang@mail.ntut.edu.tw) (H.-C. Chuang).  
<sup>1</sup> Authors contributed equally.

<https://doi.org/10.1016/j.electacta.2022.140672>  
Received 10 January 2022; Received in revised form 25 May 2022; Accepted 1 June 2022  
Available online 3 June 2022  
0013-4686/© 2022 Elsevier Ltd. All rights reserved.

Shobana Sebastin Mary Manickaraj<sup>a,b,1</sup>, Sabarison Pandiyarajan<sup>a,b,1</sup>, Ai-Ho Liao<sup>c,d</sup>, Atchaya Ramachandran<sup>e</sup>, Sheng-Tung Huang<sup>a</sup>, Priyadharshini Natarajan<sup>f</sup>, Ho-Chiao Chuang<sup>b,\*</sup>

<sup>a</sup> Department of Chemical Engineering and Biotechnology, National Taipei University of Technology, Taipei 106344, Taiwan  
<sup>b</sup> Department of Mechanical Engineering, National Taipei University of Technology, Taipei 106344, Taiwan  
<sup>c</sup> Graduate Institute of Biomedical Engineering, National Taiwan University of Science and Technology, Taipei 106335, Taiwan  
<sup>d</sup> Department of Biomedical Engineering, National Defense Medical Center, Taipei 114201, Taiwan  
<sup>e</sup> PG & Research Department of Chemistry, Bishop Heber College, Tiruchirappalli 620017, India  
<sup>f</sup> Syngene International Pvt., Ltd. Bangalore, India


作者排序 (W2)

第一作者 (X1)  
 通訊作者 (X1)  
 第二作者 (X0.8)  
 第三作者 (X0.6)  
 第四作者 (X0.4)  
 第五作者以上 (X0.2)

\* Corresponding author. (通訊作者)  
E-mail address: [hchuang@mail.ntut.edu.tw](mailto:hchuang@mail.ntut.edu.tw) (H.-C. Chuang)  
<sup>1</sup> Authors contributed equally.

# 【共通】研究彈薪申請表論文面向-B18

以林律吟教授之論文為例：  
本篇文章林教授為**第二作者**(作者順序為第二位)，對應法規應**x0.8**



Tsung-Rong Kuo<sup>1 a, b, \*</sup>, **Lu-Yin Lin<sup>2 c</sup>**, Subbiramaniyan Kubendhiran<sup>3 c</sup>, Yi-Chiun Li<sup>4</sup>, Ren-Jei Chung<sup>5</sup>, Sibidou Yougbare<sup>6 d</sup>

<sup>a</sup> International PhD Program in Biomedical Engineering, College of Biomedical Engineering, Taipei Medical University, Taipei 11031, Taiwan  
<sup>b</sup> Graduate Institute of Nanomedicine and Medical Engineering, College of Biomedical Engineering, Taipei Medical University, Taipei 11031, Taiwan  
<sup>c</sup> Department of Chemical Engineering and Biotechnology, National Taipei University of Technology, Taipei, Taiwan  
<sup>d</sup> Institut de Recherche en Sciences de la Santé/Direction Régionale du Centre Ouest (IRSS/DRCO), BP 218, 11, Nanoro, Burkina Faso


- 作者排序 (W2)
- 第一作者(x1)
  - 通訊作者(x1)
  - 第二作者(x0.8)
  - 第三作者(x0.6)
  - 第四作者(x0.4)
  - 第五作者以上(x0.2)

(通訊作者非林教授)

\* Correspondence to: T.R. Kuo, International PhD Program in Biomedical Engineering, College of Biomedical Engineering, Taipei Medical University, Taipei 11031, Taiwan.  
E-mail addresses: [trkuo@tmu.edu.tw](mailto:trkuo@tmu.edu.tw) (T.-R. Kuo), [lylin@ntut.edu.tw](mailto:lylin@ntut.edu.tw) (L.-Y. Lin), [kubechem@gmail.com](mailto:kubechem@gmail.com) (S. Kubendhiran), [piay31302@gmail.com](mailto:piay31302@gmail.com) (Y.-C. Li), [rjchung@ntut.edu.tw](mailto:rjchung@ntut.edu.tw) (R.-J. Chung), [ysibidou@gmail.com](mailto:ysibidou@gmail.com) (S. Yougbare).

# 【共通】研究彈薪申請表論文面向-B19

以林律吟教授之論文為例：  
本篇文章林教授為第三作者(作者順序為第三位)，對應法規應x0.6



Journal of Energy Storage 55 (2022) 105420

Contents lists available at ScienceDirect

Journal of Energy Storage

journal homepage: [www.elsevier.com/locate/est](http://www.elsevier.com/locate/est)

Research Papers

Investigating energy storage ability of MIL101-(Fe) derivatives prepared using successive carbonization and oxidation for supercapacitors

Yung-Fu Wu<sup>a,1</sup>, Tsung-Rong Kuo<sup>b,c,1,\*</sup>, Lu-Yin Lin<sup>d</sup>, Subbiramaniyan Kubendhiran<sup>d</sup>, Kuan-Chen Lai<sup>d</sup>, Tzu-Yang Chen<sup>d</sup>, Sibidou Yougbare<sup>e</sup>

<sup>a</sup> Department of Chemical Engineering, Ming Chi University of Technology, New Taipei City 24301, Taiwan  
<sup>b</sup> International PhD Program in Biomedical Engineering, College of Biomedical Engineering, Taipei Medical University, Taipei 11031, Taiwan  
<sup>c</sup> Graduate Institute of Nanomedicine and Medical Engineering, College of Biomedical Engineering, Taipei Medical University, Taipei 11031, Taiwan  
<sup>d</sup> Department of Chemical Engineering and Biotechnology, National Taipei University of Technology, Taipei, Taiwan  
<sup>e</sup> Institut de Recherche en Sciences de la Santé (IRSS-DRCO)/Nanoro, 03 B.P 7192, Ouagadougou 03, Burkina Faso

ARTICLE INFO

Keywords:  $\alpha$ -Fe<sub>2</sub>O<sub>3</sub>; Carbonization; Metal organic framework; MIL101(Fe); Successive carbonization/oxidation; Supercapacitor

ABSTRACT

Metal organic framework (MOF) with high surface area and tunable porous size is largely used as active material of supercapacitor (SC). MIL-101(Fe) composed of iron ions and terephthalic acid ligand is candidate active material of SC owing to its possible formation of carbon and iron compounds. Combining carbon and metal compound is feasible to establish efficient active material with ion adsorption/desorption and redox reaction charge storage abilities. In this study, it is the first time to investigate physical and electrochemical properties of MIL101(Fe) derivatives synthesized using carbonization and successive carbonization/oxidation processes as active materials of SC. Carbonization temperature of MIL-101(Fe) is optimized regarding to morphology, composition and defect/graphitization ratio. The highest specific capacitance ( $C_p$ ) of 95.7 F/g at 20 mV/s is obtained for the carbonized MIL-101(Fe) (MIL101(Fe)-C) prepared at 800 °C, due to rough surface, hollow structure and suitable defect to graphitization ratio. The MIL-101(Fe) and the successive carbonization/oxidation synthesized derivative electrodes merely achieve  $C_p$  values of 44.3 and 0.1 F/g, respectively. Symmetric SC fabricated using optimized MIL101(Fe)-C electrodes shows the maximum energy density of 1.13 Wh/kg at 400 W/kg and excellent cycling stability with  $C_p$  retention of 96% and Columbic efficiency of 72% in 8000 times repeated charging/discharging cycles.

1. Introduction

To solve serious energy shortage problems, developing efficient energy generation and storage devices are of great significance for human beings in recent years [1–8]. Batteries and capacitors are traditional energy storage devices which store charges by generating redox reactions and adsorbing/desorbing ions via static electricity, respectively [9]. By combining advantages of battery and capacitor, supercapacitor (SC) has been considered as one of effective energy storage devices owing to high specific power and long cycle life. SC stores charges by both ion adsorption/desorption and redox reactions, which mainly occurs on carbon materials and metal compounds, respectively [10,11]. Therefore, incorporating carbon materials with metal compounds as active material of SC is widely adopted to achieve excellent energy storage ability such as high specific energy and power as well as excellent cycling stability [12–14].

Metal organic framework (MOF) has been intensively applied as active material of SC, due to its high surface area and tunable porous size [15–18]. MOF with iron centers such as MIL-101(Fe) composed of coordinated iron ions linked by terephthalic acid ligands is promising active material of SC, due to the possible conversions to carbon material and iron compounds [19–21]. In previous studies, MIL-101(Fe) was commonly combined with carbon materials as active material of SC. Liu et al. prepared growth-oriented Fe-based MOF synergized with graphene aerogels composite for SC [11]. The carbonization process was also applied on MIL-101 to fabricate carbon and metal oxide composites.

\* Corresponding author at: International PhD Program in Biomedical Engineering, College of Biomedical Engineering, Taipei Medical University, Taipei 11031, Taiwan.  
E-mail address: [trkuo@tmu.edu.tw](mailto:trkuo@tmu.edu.tw) (T.-R. Kuo).  
<sup>1</sup> The authors contributed equally.

<https://doi.org/10.1016/j.est.2022.105420>  
Received 2 June 2022; Received in revised form 9 July 2022; Accepted 29 July 2022  
Available online 4 August 2022  
2352-152X/© 2022 Elsevier Ltd. All rights reserved.

① Yung-Fu Wu<sup>a,1</sup>, Tsung-Rong Kuo<sup>b,c,1,\*</sup>, ② Lu-Yin Lin<sup>d</sup>, ③ Subbiramaniyan Kubendhiran<sup>d</sup>, Kuan-Chen Lai<sup>d</sup>, Tzu-Yang Chen<sup>d</sup>, Sibidou Yougbare<sup>e</sup>

<sup>a</sup> Department of Chemical Engineering, Ming Chi University of Technology, New Taipei City 24301, Taiwan  
<sup>b</sup> International PhD Program in Biomedical Engineering, College of Biomedical Engineering, Taipei Medical University, Taipei 11031, Taiwan  
<sup>c</sup> Graduate Institute of Nanomedicine and Medical Engineering, College of Biomedical Engineering, Taipei Medical University, Taipei 11031, Taiwan  
<sup>d</sup> Department of Chemical Engineering and Biotechnology, National Taipei University of Technology, Taipei, Taiwan  
<sup>e</sup> Institut de Recherche en Sciences de la Santé (IRSS-DRCO)/Nanoro, 03 B.P 7192, Ouagadougou 03, Burkina Faso

作者排序 (W2)

- 第一作者(x1)
- 通訊作者(x1)
- 第二作者(x0.8)
- 第三作者(x0.6)
- 第四作者(x0.4)
- 第五作者以上(x0.2)

\* Corresponding author at: International PhD Program in Biomedical Engineering, College of Biomedical Engineering, Taipei Medical University, Taipei 11031, Taiwan.  
E-mail address: [trkuo@tmu.edu.tw](mailto:trkuo@tmu.edu.tw) (T.-R. Kuo).  
<sup>1</sup> The authors contributed equally.

(通訊作者非林教授)

# 【共通】研究彈薪申請表論文面向-B20

以陳生明教授之論文為例：

本篇文章陳教授為**第四作者**(作者順序為第四位)，對應法規應**x0.4**

Contents lists available at ScienceDirect  
Journal of Alloys and Compounds  
journal homepage: [www.elsevier.com/locate/jalcom](http://www.elsevier.com/locate/jalcom)

Research Article  
Direct Z-scheme WO<sub>3</sub>/In<sub>2</sub>S<sub>3</sub> heterostructures for enhanced photocatalytic reduction Cr(VI)

Yuxiang Hua<sup>a</sup>, Chengyao Hu<sup>b</sup>, Muhammad Arif<sup>a</sup>, Shen-ming Chen<sup>c</sup>, Min Zhang<sup>a,d,\*</sup>, Xiaoheng Liu<sup>a,\*\*</sup>

<sup>a</sup> Key Laboratory of Education Ministry for Soft Chemistry and Functional Materials, Nanjing University of Science and Technology, Nanjing 210094, China  
<sup>b</sup> Advanced Analysis and Testing Center, Nanjing Forestry University, Nanjing 210037, China  
<sup>c</sup> Department of Chemical Engineering and Biotechnology, National Taipei University of Technology, Taipei 106, Taiwan, ROC  
<sup>d</sup> College of Chemistry and Material Engineering, Quzhou University, Quzhou, Zhejiang 324000, China

ARTICLE INFO  
Article history:  
Received 24 May 2021  
Received in revised form 2 March 2022  
Accepted 7 March 2022  
Available online 11 March 2022

Keywords:  
In<sub>2</sub>S<sub>3</sub>  
WO<sub>3</sub>  
Z-scheme photocatalyst  
Photocatalytic Cr(VI) reduction

1. Introduction  
With the development of industry, potentially toxic metals ions pose a major threat to the water environment [1–3]. As a common metal-chromium ions, it is widely used in electroplating, leather tanning, steelmaking, and chemical manufacturing [4–6]. Since Cr(VI) has a regular tetrahedral structure similar to PO<sub>4</sub><sup>3-</sup> and SO<sub>4</sub><sup>2-</sup>, it can easily enter cells through anion channels, which seriously affects human health and safety [7]. In 2019, chromium compounds with hexavalent were included in the list of toxic and harmful water pollutants. As we all know, Cr(III) as a trace element of the human body, has the advantages of low mobility in aquatic environment and easy formation of Cr(OH)<sub>3</sub> precipitation in neutral or alkaline environments, which has become an effective way

ABSTRACT  
The design of efficient and stable photocatalysts for the removal of heavy metals in the environment has become a research hotspot. Here, a composite photocatalyst with three-dimensional In<sub>2</sub>S<sub>3</sub> microspheres supported by WO<sub>3</sub> nanoparticles was synthesized for the photoreduction of Cr(VI) for the first time. The constructed composite catalyst has a direct Z-scheme electron transport mechanism without any precious metals (Au, Pt, and Ag), quantum dots (TiO<sub>2</sub> QDs) or carbon materials (Graphene) as electronic media. Constructing a direct Z-scheme WO<sub>3</sub>/In<sub>2</sub>S<sub>3</sub> photocatalyst can greatly retain the reduction and oxidation reaction sites on the surface of the heterojunction and accelerate the reduction reaction. Under visible light irradiation, it greatly promotes the photocatalytic reduction of Cr(VI), which is 67.7 times and 3.6 times the reduction rates of WO<sub>3</sub> and In<sub>2</sub>S<sub>3</sub>, respectively. The favorable photocatalytic performance of WO<sub>3</sub>/In<sub>2</sub>S<sub>3</sub> should be attributed to the effective interfacial contact between the semiconductors in the Z-scheme system, thereby realizing effective electron transfer and charge separation. In addition, the stability of WO<sub>3</sub>/In<sub>2</sub>S<sub>3</sub> was studied, and a possible mechanism in the photoreduction process of Cr(VI) was proposed.  
© 2022 Elsevier B.V. All rights reserved.

to solve the pollution of Cr(VI) [8]. Generally, the sulfite or ferrous salt was used in industry to reduce Cr(VI) to Cr(III) in an acidic environment, and then alkali treatment is performed to obtain precipitate [9]. This method was prone to produce secondary solid waste and SO<sub>2</sub>, which poses environmental hazards. In recent years, semiconductor photocatalysts have been generated electron-hole pairs under light excitation, in which electrons have strong reducibility without any pollution, and can be used to reduce Cr(VI) [10]. Wang et al. prepared CeO<sub>2</sub> nanotubes by a surfactant-assisted hydrothermal method for photoreduction of Cr(VI). The pure CeO<sub>2</sub> has weak photoreduction performance without adding oxalic acid [11]. Ren's group reported the use of ZnO to reduce Cr(VI) under ultraviolet light [12].  
The photocatalysis has become a "green technology" for addressing environmental problems. To achieve the goal of photoreduction of Cr(VI), photocatalysts need to have narrower band gaps, more negative conduction band (CB) sites, and more positive valence band (VB) sites. This is difficult for a single photocatalytic material to have at the same time. The heterojunction catalytic system with Z-scheme electron transport mechanism can not only decrease the photo-generated electron-hole recombination rate, but also retain

\* Corresponding author at: Key Laboratory of Education Ministry for Soft Chemistry and Functional Materials, Nanjing University of Science and Technology, Nanjing 210094, China  
E-mail addresses: [mzhang.chemmater@qzc.edu.cn](mailto:mzhang.chemmater@qzc.edu.cn) (M. Zhang), [xhliu@njjust.edu.cn](mailto:xhliu@njjust.edu.cn) (X. Liu).

https://doi.org/10.1016/j.jalcom.2022.164488  
0925-8388/© 2022 Elsevier B.V. All rights reserved.

Yuxiang Hua<sup>a</sup>, Chengyao Hu<sup>b</sup>, Muhammad Arif<sup>a</sup>, **Shen-ming Chen<sup>c</sup>**, Min Zhang<sup>a,d,\*</sup>, Xiaoheng Liu<sup>a,\*\*</sup>

<sup>a</sup> Key Laboratory of Education Ministry for Soft Chemistry and Functional Materials, Nanjing University of Science and Technology, Nanjing 210094, China  
<sup>b</sup> Advanced Analysis and Testing Center, Nanjing Forestry University, Nanjing 210037, China  
<sup>c</sup> Department of Chemical Engineering and Biotechnology, National Taipei University of Technology, Taipei 106, Taiwan, ROC  
<sup>d</sup> College of Chemistry and Material Engineering, Quzhou University, Quzhou, Zhejiang 324000, China

\* Corresponding author at: Key Laboratory of Education Ministry for Soft Chemistry and Functional Materials, Nanjing University of Science and Technology, Nanjing 210094, China  
\*\* **Corresponding author.** (通訊作者非陳教授)  
E-mail addresses: [mzhang.chemmater@qzc.edu.cn](mailto:mzhang.chemmater@qzc.edu.cn) (M. Zhang), [xhliu@njjust.edu.cn](mailto:xhliu@njjust.edu.cn) (X. Liu).

作者排序 (W2)

- 第一作者(x1)
- 通訊作者(x1)
- 第二作者(x0.8)
- 第三作者(x0.6)
- 第四作者(x0.4)
- 第五作者以上(x0.2)

# 【共通】研究彈薪申請表論文面向-B21

以莊賀喬教授之論文為例：  
本篇文章莊教授為第五作者(作者順序為第五位)，對應法規應x0.2

Received: 19 May 2022 | Revised: 29 October 2022 | Accepted: 8 November 2022  
DOI: 10.1002/btm2.10450

RESEARCH ARTICLE

Combined use of microbubbles of various sizes and single-transducer dual-frequency ultrasound for safe and efficient inner ear drug delivery

Ai-Ho Liao<sup>1,2</sup> | Chih-Hung Wang<sup>3,4</sup> | Bo-Han Wang<sup>5</sup> | Yi-Chun Lin<sup>4</sup> | **Ho-Chiao Chuang<sup>5</sup>** | Hao-Li Liu<sup>6</sup> | Cheng-Ping Shih<sup>3</sup>

<sup>1</sup>Graduate Institute of Biomedical Engineering, National Taiwan University of Science and Technology, Taipei, Taiwan  
<sup>2</sup>Department of Biomedical Engineering, National Defense Medical Center, Taipei, Taiwan  
<sup>3</sup>Department of Otolaryngology-Head and Neck Surgery, Tri-Service General Hospital, National Defense Medical Center, Taipei, Taiwan  
<sup>4</sup>Graduate Institute of Medical Sciences, National Defense Medical Center, Taipei, Taiwan  
<sup>5</sup>Department of Mechanical Engineering, National Taipei University of Technology, Taipei, Taiwan  
<sup>6</sup>Department of Electrical Engineering, National Taiwan University, Taipei, Taiwan

**Correspondence**  
Cheng-Ping Shih, Department of Otolaryngology-Head and Neck Surgery, Tri-Service General Hospital, National Defense Medical Center, No. 325, Sec. 2, Chenggong Rd., Neihu District, Taipei City 11490, Taiwan  
Email: eric660802@gmail.com, zhengping\_shi@yahoo.com.tw

**Funding information**  
Ministry of Science and Technology, Taiwan. Grant/Award Numbers: MOST108-2311-B-011-161-MY4, MOST110-2311-B-016-047-MY3, MOST110-2311-B-016-045, MOST109-2311-B-016-005, National Taiwan University of Science and Technology. Grant/Award Numbers: NTUST-TSGH-110-01, NTUST-TSGH-111-01; Tri-Service General Hospital. Grant/Award Numbers: TSGH-D-111059, TSGH-A-111001, TSGH-A-110001; Medical Affairs Bureau, Ministry of National Defense, ROC. Grant/Award Number: MAB-D-111145

**Abstract**  
We have previously applied ultrasound (US) with microbubbles (MBs) to enhance inner ear drug delivery, with most experiments conducted using single-frequency, high-power density US, and multiple treatments. In the present study, the treatment efficacy was enhanced and safety concerns were addressed using a combination of low-power-density, single-transducer, dual-frequency US ( $I_{SPTA} = 213 \text{ mW/cm}^2$ ) and MBs of different sizes coated with insulin-like growth factor-1 (IGF-1). This study is the first to investigate the drug-coating capacity of human serum albumin (HSA) MBs of different particle sizes and their drug delivery efficiency. The concentration of HSA was adjusted to produce different MB sizes. The drug-coating efficiency was significantly higher for large-sized MBs than for smaller MBs. In vitro Franz diffusion experiments showed that the combination of dual-frequency US and large MB size delivered the most IGF-1 ( $24.3 \pm 0.47 \text{ ng/cm}^2$ ) to the receptor side at the second hour of treatment. In an in vivo guinea pig experiment, the efficiency of IGF-1 delivery into the inner ear was 15.9 times greater in animals treated with the combination of dual-frequency US and large MBs (D-USMB) than in control animals treated with round window soaking (RWS). The IGF-1 delivery efficiency was 10.15 times greater with the combination of single-frequency US and large size MBs (S-USMB) than with RWS. Confocal microscopy of the cochlea showed a stronger distribution of IGF-1 in the basal turn in the D-USMB and S-USMB groups than in the RWS group. In the second and third turns, the D-USMB group showed the greatest IGF-1 distribution.

Ai-Ho Liao and Chih-Hung Wang contributed equally to this study.

This is an open access article under the terms of the [Creative Commons Attribution License](#), which permits use, distribution and reproduction in any medium, provided the original work is properly cited.  
© 2022 The Authors. Bioengineering & Translational Medicine published by Wiley Periodicals LLC on behalf of American Institute of Chemical Engineers.

Bioeng Transl Med. 2023;8:e10450.  
<https://doi.org/10.1002/btm2.10450> | 1 of 14

Ai-Ho Liao<sup>1,2</sup> | Chih-Hung Wang<sup>3,4</sup> | Bo-Han Wang<sup>5</sup> | Yi-Chun Lin<sup>4</sup> | **Ho-Chiao Chuang<sup>5</sup>** | Hao-Li Liu<sup>6</sup> | Cheng-Ping Shih<sup>3</sup>

<sup>1</sup>Graduate Institute of Biomedical Engineering, National Taiwan University of Science and Technology, Taipei, Taiwan  
<sup>2</sup>Department of Biomedical Engineering, National Defense Medical Center, Taipei, Taiwan  
<sup>3</sup>Department of Otolaryngology-Head and Neck Surgery, Tri-Service General Hospital, National Defense Medical Center, Taipei, Taiwan  
<sup>4</sup>Graduate Institute of Medical Sciences, National Defense Medical Center, Taipei, Taiwan  
<sup>5</sup>Department of Mechanical Engineering, National Taipei University of Technology, Taipei, Taiwan  
<sup>6</sup>Department of Electrical Engineering, National Taiwan University, Taipei, Taiwan

**Correspondence** (通訊作者非莊教授)  
Cheng-Ping Shih, Department of Otolaryngology-Head and Neck Surgery, Tri-Service General Hospital, National Defense Medical Center, No. 325, Sec. 2, Chenggong Rd., Neihu District, Taipei City 11490, Taiwan.  
Email: eric660802@gmail.com, zhengping\_shi@yahoo.com.tw

作者排序 (W2)

- 第一作者(x1)
- 通訊作者(x1)
- 第二作者(x0.8)
- 第三作者(x0.6)
- 第四作者(x0.4)
- 第五作者以上 (x0.2)

# 【共通】研究彈薪申請表論文面向-B22

查詢W2方式

注意事項:

1. 需檢附論文第一頁為佐證資料。
2. 第一作者到第五作者依由左至右的順序排列。
3. 通訊作者通常會使用\*字號、或是下方會有一欄 **Corresponding author** 的部份，如還查不到的話建議使用WOS查詢。

WOS查詢畫面:

**An Upper Extremity Rehabilitation System Using Efficient Vision-Based Action Identification Techniques**

By Chen, YL (Chen, Yen-Lin) [1]; Liu, CH (Liu, Chin-Hsuan) [1]; Yu, CW (Yu, Chao-Wei) [1]; Lee, P (Lee, Posen) [2]; Kuo, YW (Kuo, Yao-Wen) [1]

View Web of Science ResearcherID and ORCID (provided by Clarivate)

Source APPLIED SCIENCES-BASEL  
Volume: 8 Issue: 7  
DOI: 10.3390/app8071161

Article Number 1161

Published JUL 2018

Indexed 2018-09-07

Document Type Article

Abstract This study proposes an action identification system for home upper extremity rehabilitation. In the proposed system, we apply an RGB-depth (color-depth) sensor to capture the image sequences of the patient's upper extremity actions to identify its movements. We apply a skin color detection technique to assist with extremity identification and to build up the upper extremity skeleton points. We use the dynamic time warping algorithm to determine the rehabilitation actions. The system presented herein builds up upper extremity skeleton points rapidly. Through the upper extremity of the human skeleton and human skin color information, the upper extremity skeleton points are effectively established by the proposed system, and the rehabilitation actions of patients are identified by a dynamic time warping algorithm. Thus, the proposed system can achieve a high recognition rate of 98% for the defined rehabilitation actions for the various muscles. Moreover, the computational speed of the proposed system can reach 125 frames per second-the processing time per frame is less than 8 ms on a personal computer platform. This computational efficiency allows efficient extensibility for future developments to deal with complex ambient environments and for implementation in embedded and pervasive systems. The major contributions of the study are: (1) the proposed system is not only a physical exercise game, but also a movement training program for specific muscle groups; (2) The hardware of upper extremity rehabilitation system included a personal computer with personal computer and a depth camera. These are economic equipment, so that patients who need this system can set up one set at home; (3) patients can perform rehabilitation actions in sitting position to prevent him/her from falling down during training; (4) the accuracy rate of identifying rehabilitation action is as high as 98%, which is sufficient for distinguishing between correct and wrong action when performing specific action trainings; (5) The proposed upper extremity rehabilitation system is real-time, efficient to vision-based action identification, and low-cost hardware and software, which is affordable for most families.

Keywords Author Keywords: upper extremity identification; color and depth sensors; skeleton points; rehabilitation actions; home rehabilitation; computer vision  
Keywords Plus: COST-EFFECTIVENESS; TELEMEDICINE; CARE; BALANCE; TELEHEALTH; TOOL

Author Information Corresponding Address: Lee, Posen (corresponding author)  
I Shou Univ, Dept Occupat Therapy, Kaohsiung 82445, Taiwan

Author Information

Corresponding Address: Lee, Posen (corresponding author)

▲ I Shou Univ, Dept Occupat Therapy, Kaohsiung 82445, Taiwan

Author Information

Corresponding Address: Lee, Posen (corresponding author)

▲ I Shou Univ, Dept Occupat Therapy, Kaohsiung 82445, Taiwan

續下頁



# 【共通】研究彈薪申請表論文面向-B23

查詢W3方式

依ppt第30.31頁方式下載論文檔案

以莊賀喬教授之論文為例: (只有一位通訊作者之狀況)

本篇文章通訊作者只有一位，對應法規應x1

Electrochimica Acta  
journal homepage: [www.journals.elsevier.com/electrochimica-acta](http://www.journals.elsevier.com/electrochimica-acta)

*Sansevieria trifasciata* biomass-derived activated carbon by supercritical-CO<sub>2</sub> route: Electrochemical detection towards carcinogenic organic pollutant and energy storage application

Shobana Sebastin Mary Manickaraj<sup>a,b,1</sup>, Sabarison Pandiyarajan<sup>a,b,1</sup>, Ai-Ho Liao<sup>c,d</sup>, Atchaya Ramachandran<sup>e</sup>, Sheng-Tung Huang<sup>a</sup>, Priyadharshini Natarajan<sup>f</sup>, Ho-Chiao Chuang<sup>b,\*</sup>

<sup>a</sup> Department of Chemical Engineering and Biotechnology, National Taipei University of Technology, Taipei 106344, Taiwan  
<sup>b</sup> Department of Mechanical Engineering, National Taipei University of Technology, Taipei 106344, Taiwan  
<sup>c</sup> Graduate Institute of Biomedical Engineering, National Taiwan University of Science and Technology, Taipei 106335, Taiwan  
<sup>d</sup> Department of Biomedical Engineering, National Defense Medical Center, Taipei 114201, Taiwan  
<sup>e</sup> PG & Research Department of Chemistry, Bishop Heber College, Tiruchirappalli 620017, India  
<sup>f</sup> Syngene International Pvt., Ltd. Bangalore, India

**ARTICLE INFO**

**Keywords:**  
Activated carbon  
Sansevieria trifasciata  
Supercritical-CO<sub>2</sub>  
Electrochemical sensor  
Supercapacitor

**ABSTRACT**

Activated carbon (AC) has been widely used for electrochemical applications, such as electrochemical sensors, energy storage applications, etc., due to its fine porous structure, volumetric capacitance, and chemical stability. Supercritical-CO<sub>2</sub> (SC-CO<sub>2</sub>) has a fascinating advantage in material science due to its microbubble cavitation, high diffusivity, and high permeability. In the shed of light, we developed a high porous *Sansevieria trifasciata* biomass-derived AC by SC-CO<sub>2</sub> (SC-ST-AC). For comparison purposes, the AC was also prepared in a conventional approach (C-ST-AC). The prepared ACs were characterized through various spectroscopic and microscopic techniques to study their surface morphological character, structural analysis, and phase purity. The electrochemical performance was evaluated by two different applications: electrochemical detection and energy storage application. Based on the results, the SC-ST-AC exhibits higher porous architecture in their morphology and high phase purity with amorphous nature than C-ST-AC. In the preliminary electrochemical analysis, SC-ST-AC achieved higher performance than C-ST-AC. Thus, SC-ST-AC is applied to the real-time application and it exposed a superior limit of detection (0.005 μM L<sup>-1</sup>) and sensitivity (0.854 μA μM<sup>-1</sup> cm<sup>-2</sup>) towards MA sensing and higher specific capacitance (342.5 F/g for 2 A/g) with 92.09 % of retention at high current density. Thereby, we suggest the SC-CO<sub>2</sub> method is a promising approach to develop a highly porous carbon material with excellent electrochemical performance.

**1. Introduction**

In recent eras, carbon-based materials including one-dimension (1D) carbon nanotubes, carbon nanofibers [1,2], two-dimension (2D) graphene [3], three-dimension (3D) graphite, activated carbon, and its derivatives [4,5] have been extensively investigated as successful commercialization materials in several sectors. Among them activated carbon (AC) is considered the most cardinal material for electrochemical application owing to its high surface area, porous architecture, and chemical stability [6-8]. The varieties of functional group moiety fascinated on the surface make it as a promising electrode material for energy storage applications [9]. Traditionally, the preparation of AC is done by the pyrolysis of fossil raw materials such as coal and petroleum coke or wood, followed by a physical or chemical activation process [10]. Due to the rapid increase of the global population and economy, the demand for energy and resources is also increasing exponentially, resulting in a lack of fossil fuels [11]. Therefore, cost-effective renewable carbon sources, the development of economic efficiency methods, and environmental safety are all issues that must be thoroughly investigated to produce advanced activated carbon that is more environmentally friendly. In this regard, biomass materials are presently recognized as the most viable candidates for preparing carbon materials

\* Corresponding author.  
E-mail address: [hchuang@mail.ntut.edu.tw](mailto:hchuang@mail.ntut.edu.tw) (H.-C. Chuang).  
<sup>1</sup> Authors contributed equally.

<https://doi.org/10.1016/j.electacta.2022.140672>  
 Received 10 January 2022; Received in revised form 25 May 2022; Accepted 1 June 2022  
 Available online 3 June 2022  
 0013-4686/© 2022 Elsevier Ltd. All rights reserved.

Shobana Sebastin Mary Manickaraj<sup>a,b,1</sup>, Sabarison Pandiyarajan<sup>a,b,1</sup>, Ai-Ho Liao<sup>c,d</sup>, Atchaya Ramachandran<sup>e</sup>, Sheng-Tung Huang<sup>a</sup>, Priyadharshini Natarajan<sup>f</sup>, Ho-Chiao Chuang<sup>b,\*</sup>

<sup>a</sup> Department of Chemical Engineering and Biotechnology, National Taipei University of Technology, Taipei 106344, Taiwan  
<sup>b</sup> Department of Mechanical Engineering, National Taipei University of Technology, Taipei 106344, Taiwan  
<sup>c</sup> Graduate Institute of Biomedical Engineering, National Taiwan University of Science and Technology, Taipei 106335, Taiwan  
<sup>d</sup> Department of Biomedical Engineering, National Defense Medical Center, Taipei 114201, Taiwan  
<sup>e</sup> PG & Research Department of Chemistry, Bishop Heber College, Tiruchirappalli 620017, India  
<sup>f</sup> Syngene International Pvt., Ltd. Bangalore, India

共同作者數 (W3)

無(x1)

1位通訊作者(x1)

2位(含)以上通訊作者(x0.8)

有多位作者 Equal Contribution (x0.8)

\* Corresponding author. (通訊作者)  
 E-mail address: [hchuang@mail.ntut.edu.tw](mailto:hchuang@mail.ntut.edu.tw) (H.-C. Chuang)  
<sup>1</sup> Authors contributed equally.

# 【共通】研究彈薪申請表論文面向-B24

以莊賀喬教授之論文為例: (有兩位通訊作者之狀況)  
 本篇文章通訊作者有兩位且均為國內學者, 對應法規應x0.8

Contents lists available at ScienceDirect  
 Biomedical Signal Processing and Control  
 journal homepage: [www.elsevier.com/locate/bspc](http://www.elsevier.com/locate/bspc)

Combining the wavelet transform with a phase-lead compensator to a respiratory motion compensation system with an ultrasound tracking technique in radiation therapy

Chia-Chun Kuo<sup>a,d,e</sup>, Ming-Lu Guo<sup>g</sup>, Ai-Ho Liao<sup>b,c</sup>, Hsiao-Wei Yu<sup>a,f</sup>, Muthusankar Ganesan<sup>g,k</sup>, Chu-Wei Li<sup>g</sup>, Shiu-Chen Jeng<sup>a,h</sup>, Jeng-Fong Chiou<sup>a,i,j</sup>, Lai-Lei Ting<sup>a,\*</sup>, Ho-Chiao Chuang<sup>g,\*</sup>

<sup>a</sup> Department of Radiation Oncology, Taipei Medical University Hospital, Taipei, Taiwan  
<sup>b</sup> Graduate Institute of Biomedical Engineering, National Taiwan University of Science and Technology, Taipei, Taiwan  
<sup>c</sup> Department of Biomedical Engineering, National Defense Medical Center, Taipei, Taiwan  
<sup>d</sup> Department of Radiation Oncology, Wan Fang Hospital, Taipei Medical University, Taipei, Taiwan  
<sup>e</sup> School of Health Care Administration, College of Management, Taipei Medical University, Taipei, Taiwan  
<sup>f</sup> School of Biomedical Engineering, College of Biomedical Engineering, Taipei Medical University, Taipei, Taiwan  
<sup>g</sup> Department of Mechanical Engineering, National Taipei University of Technology, Taipei, Taiwan  
<sup>h</sup> School of Dentistry, College of Oral Medicine, Taipei Medical University, Taipei, Taiwan  
<sup>i</sup> Department of Radiology, School of Medicine, College of Medicine, Taipei Medical University, Taipei, Taiwan  
<sup>j</sup> Taipei Cancer Center, Taipei Medical University, Taipei, Taiwan  
<sup>k</sup> Department of Industrial Chemistry, Alagappa University, Karaikudi 630001, Tamil Nadu, India

ARTICLE INFO  
 Keywords: Wavelet transform, Respiratory motion compensation, Ultrasound image tracking

ABSTRACT  
 This study evaluated the feasibility of applying the wavelet transform (WT) combined with a phase-lead compensator (PLC) to our previously developed two-dimensional respiratory motion compensation system (RMCS). This system automatically and instantaneously adjusts PLC parameters according to different respiration signals to reduce influences of the system delay time, improving the compensation effect of the RMCS during respiratory motion compensation. This study performed respiratory movement compensation experiments with a two-dimensional respiratory motion simulation system (RMSS) and the RMCS. Human respiratory signals were captured using our previously developed ultrasound image tracking algorithm (UITA). In this study, a displacement compensation RMCS algorithm based on the combination of WT and PLC was developed by LabVIEW, which allows an automatic adjustment of the PLC parameters according to various respiratory waveforms, achieving a better compensation effect. The experiment results indicated that the compensation rate (CR) of right-left and superior-inferior directions had both improved 67.96-88.05% and 70.38-91.43%, respectively. In this study, the proposed method combined with WT and PLC applied in respiratory movement compensation experiments; the UITA was used for tracking diaphragm motion which substitutes for tumor motion. This noninvasive monitoring method also helps reduce side effects after treatment. The experimental results indicated that the effect of using the WT combined with the PLC to compensate for various respiratory signals was improved over our previously developed compensation algorithm.

1. Introduction  
 During radiotherapy, the anatomical structure and location of a lesion are usually different from those of the target used in the treatment planning system. One of the main reasons for this is the organ movement that occurs while breathing, which also causes the tumor to deviate from the original irradiation target position during the treatment [1-3]. The tumor movement makes actual dose distribution differ from the expected dose distribution, resulting insufficient dose coverage on target tumor and excessive dose on surrounding tissues. The unwanted dose distribution increases serious side effects and great reduction of treatment effectiveness. Langen et al. [4] documented many types of organ movements, including types of the liver, diaphragm, kidney, pancreas, lung tumors, and prostate. Diaphragm and liver are affected by

\* Corresponding authors.  
 E-mail addresses: [llt1957@gmail.com](mailto:llt1957@gmail.com) (L.-L. Ting), [hchuang@mail.ntut.edu.tw](mailto:hchuang@mail.ntut.edu.tw) (H.-C. Chuang).

<https://doi.org/10.1016/j.bspc.2022.103892>  
 Received 21 March 2022; Received in revised form 23 May 2022; Accepted 12 June 2022  
 Available online 17 June 2022  
 1746-8094/© 2022 Elsevier Ltd. All rights reserved.

Chia-Chun Kuo<sup>a,d,e</sup>, Ming-Lu Guo<sup>g</sup>, Ai-Ho Liao<sup>b,c</sup>, Hsiao-Wei Yu<sup>a,f</sup>, Muthusankar Ganesan<sup>g,k</sup>, Chu-Wei Li<sup>g</sup>, Shiu-Chen Jeng<sup>a,h</sup>, Jeng-Fong Chiou<sup>a,i,j</sup>, Lai-Lei Ting<sup>a,\*</sup>, Ho-Chiao Chuang<sup>g,\*</sup>

<sup>a</sup> Department of Radiation Oncology, Taipei Medical University Hospital, Taipei, Taiwan  
<sup>b</sup> Graduate Institute of Biomedical Engineering, National Taiwan University of Science and Technology, Taipei, Taiwan  
<sup>c</sup> Department of Biomedical Engineering, National Defense Medical Center, Taipei, Taiwan  
<sup>d</sup> Department of Radiation Oncology, Wan Fang Hospital, Taipei Medical University, Taipei, Taiwan  
<sup>e</sup> School of Health Care Administration, College of Management, Taipei Medical University, Taipei, Taiwan  
<sup>f</sup> School of Biomedical Engineering, College of Biomedical Engineering, Taipei Medical University, Taipei, Taiwan  
<sup>g</sup> Department of Mechanical Engineering, National Taipei University of Technology, Taipei, Taiwan  
<sup>h</sup> School of Dentistry, College of Oral Medicine, Taipei Medical University, Taipei, Taiwan  
<sup>i</sup> Department of Radiology, School of Medicine, College of Medicine, Taipei Medical University, Taipei, Taiwan  
<sup>j</sup> Taipei Cancer Center, Taipei Medical University, Taipei, Taiwan  
<sup>k</sup> Department of Industrial Chemistry, Alagappa University, Karaikudi 630001, Tamil Nadu, India

共同作者數 (W3)

無(x1)  
 1位通訊作者(x1)  
 2位(含)以上通訊作者(x0.8)  
 有多位作者 Equal Contribution (x0.8)

(通訊作者)  
 \* Corresponding authors.  
 E-mail addresses: [llt1957@gmail.com](mailto:llt1957@gmail.com) (L.-L. Ting), [hchuang@mail.ntut.edu.tw](mailto:hchuang@mail.ntut.edu.tw) (H.-C. Chuang).

# 【共通】研究彈薪申請表論文面向-B25

以莊賀喬教授之論文為例：(有兩位通訊作者之狀況)  
本篇文章通訊作者有兩位，除莊教授外另外一位為國外學者，對應法規應x1

Contents lists available at ScienceDirect  
Journal of Alloys and Compounds  
journal homepage: www.elsevier.com/locate/jalcom

Robust fabrication of silver pyro-vanadates via sonochemical approach for advanced energy storage application

Sabarison Pandiyarajan<sup>a,b</sup>, Rajkumar Srinivasan<sup>c</sup>, Shobana Sebastin Mary Manickaraj<sup>a,b</sup>, Ho-Chiao Chuang<sup>b,\*</sup>, Princy Merlin Johnson<sup>c,\*</sup>

<sup>a</sup> Department of Chemical Engineering, National Taipei University of Technology, Taipei 10608, Taiwan  
<sup>b</sup> Department of Mechanical Engineering, National Taipei University of Technology, Taipei 10608, Taiwan  
<sup>c</sup> PG and Research Department of Chemistry, Bishop Heber College, Bharathidasan University, Tiruchirappalli, India

ARTICLE INFO

Article history:  
Received 14 June 2021  
Received in revised form 9 September 2021  
Accepted 5 October 2021  
Available online 8 October 2021

Keywords:  
Silver pyro-vanadates  
Ultrasonication, Specific capacity  
Energy storage systems

ABSTRACT

One of the major challenges in the twenty-first century is the development of ultrahigh performance electrical energy storage (ES) devices with faster, safer, and more efficient ES materials. Herein, we report newly designed silver vanadates ( $Ag_4V_2O_9$ ), which serve as significant electrode material for upcoming ES devices due to its greater electrical conductivity as well as electrochemical activity.  $Ag_4V_2O_9$  were synthesized by the ultrasonication method. The as-synthesized material was characterized with various spectral as well as analytical methods. Furthermore, the supercapacitive property of  $Ag_4V_2O_9$  was evaluated using different electroanalytical techniques. The  $Ag_4V_2O_9$  electrode exhibited well electrochemical performance with a specific capacity ( $C_{sp}$ ) of  $548 \text{ C g}^{-1}$  at the current density of  $1 \text{ Ag}^{-1}$  and significant capacity retention of 88.7% even after 5000 GCD cycles at  $6 \text{ Ag}^{-1}$ . The lowest value of charge transfer resistance ( $R_{ct} = 4.12 \text{ }\Omega$ ), and equivalent series resistance (ESR =  $6.33 \text{ }\Omega$ ) exposed the faster reaction kinetics. The superior electrochemical performance was ascribed to its unique structure, which contributes to high conductivity, easy electron transfer, short ion diffusion distances, fast kinetics as well as a huge number of active sites in the electrode material. The electrochemical results demonstrated that  $Ag_4V_2O_9$  could be utilized as electro-active material for advanced energy storage systems.

© 2021 Elsevier B.V. All rights reserved.

1. Introduction

Energy crisis is one of the most pressing problems in the current scenario. Considerations about greenhouse effect have prompted researchers to perform a detailed investigation on energy conversion as well as storage technology [1]. In order to solve this issue, fuel cells, batteries and supercapacitors have become more popular as strong candidates [2]. Supercapacitors (SCs) have received a lot of attention as a type of high-efficiency energy storage device because they can deliver more power density with a longer cycling lifespan than batteries and store more energy density than conventional capacitors. Furthermore, due to their rapid rechargeability, much greater cycling stability, and higher rate capability, SCs are good alternatives for a battery replacement if their energy density is significantly high [3–6].

Based on the principle of charge storage process, there are three types of SCs: the electric double layer (EDLC) [7], pseudocapacitors (PCs) [8] as well as hybrid capacitors [9]. The former is distinguished primarily via ion as well as electron separation at the electrode/electrolyte interface, while a Faradaic charge transfer reaction takes place at the active material in a redox pseudo capacitor. Hybrid capacitors are operating by the combination of Faradaic as well as Non-Faradaic reactions. Many researchers have made great efforts to study PCs because their energy density is substantially higher than EDLCs [10,11].

Because of the large  $C_{sp}$  and superlative redox activity, transition metal oxides (TMOs) have been found to be promising as electrodes for PCs over the last few decades [12–14]. Several TMOs, like  $RuO_2$ ,  $MnO_2$ ,  $NiO$ ,  $Co_3O_4$ ,  $MoO_3$ , and  $SnO_2$  were efficiently used as electrode materials in PCs. During the charge/discharge processes, PCs with these kinds of electrodes invariably exhibited poor stability, high resistance as well as large volume changes [15]. To address this concern, mixed TMOs have emerged as promising electrodes for SCs owing to their ability to improve electrochemical performance in terms of power stability, specific capacity as well as electrical conductivity [16]. Among the TMOs, mixed metal oxides, binary,

\* Corresponding authors.  
E-mail addresses: [sabarison6@gmail.com](mailto:sabarison6@gmail.com) (S. Pandiyarajan), [hchuang@mail.ntut.edu.tw](mailto:hchuang@mail.ntut.edu.tw) (H.-C. Chuang), [pmej\\_68@yahoo.co.in](mailto:pmej_68@yahoo.co.in) (P.M. Johnson).

<https://doi.org/10.1016/j.jalcom.2021.162268>  
0925-8388/© 2021 Elsevier B.V. All rights reserved.

Sabarison Pandiyarajan<sup>a,b</sup>, Rajkumar Srinivasan<sup>c</sup>, Shobana Sebastin Mary Manickaraj<sup>a,b</sup>,  
Ho-Chiao Chuang<sup>b,\*</sup>, Princy Merlin Johnson<sup>c,\*</sup>

<sup>a</sup> Department of Chemical Engineering, National Taipei University of Technology, Taipei 10608, Taiwan  
<sup>b</sup> Department of Mechanical Engineering, National Taipei University of Technology, Taipei 10608, Taiwan  
<sup>c</sup> PG and Research Department of Chemistry, Bishop Heber College, Bharathidasan University, Tiruchirappalli, India

共同作者數 (W3)

無 (x1)

1位通訊作者 (x1)

2位(含)以上通訊作者 (x0.8)

有多位作者 Equal Contribution (x0.8)

(通訊作者)

\* Corresponding authors.  
E-mail addresses: [sabarison6@gmail.com](mailto:sabarison6@gmail.com) (S. Pandiyarajan), [hchuang@mail.ntut.edu.tw](mailto:hchuang@mail.ntut.edu.tw) (H.-C. Chuang), [pmej\\_68@yahoo.co.in](mailto:pmej_68@yahoo.co.in) (P.M. Johnson).

# 【共通】研究彈薪申請表論文面向-B26

以林律吟教授之論文為例: (有多位共同貢獻作者之狀況)  
本篇文章有**多位Equal Contribution**，對應法規應**x0.8**

Contents lists available at ScienceDirect

Journal of Alloys and Compounds

journal homepage: [www.elsevier.com/locate/jalcom](http://www.elsevier.com/locate/jalcom)

ELSEVIER

Journal of Alloys and Compounds

Investigating energy storage ability of ZIF67-derived perovskite fluoride via tuning ammonium fluoride amounts

Pin-Yan Lee<sup>a,1</sup>, Lu-Yin Lin<sup>a,b,\*,1</sup>

<sup>a</sup> Department of Chemical Engineering and Biotechnology, National Taipei University of Technology, Taipei, Taiwan  
<sup>b</sup> Research Center of Energy Conservation for New Generation of Residential, Commercial, and Industrial Sectors, Taipei, Taiwan

ARTICLE INFO

Article history:  
Received 7 July 2021  
Received in revised form 25 September 2021  
Accepted 28 September 2021  
Available online 29 September 2021

Keywords:  
Ammonia fluoride  
Ligand  
Perovskite  
Supercapacitor  
ZIF67  
2-methylimidazole

ABSTRACT

Zeolitic imidazolate framework 67 (ZIF67) is widely considered as potential active material for supercapacitors (SC) due to large surface area and tunable structures, but small electrical conductivity limits its energy storage ability. Fluoride with high electrical conductivity is reported to be beneficial on reducing charge-transfer resistance of SC. In this study, ZIF67-derived perovskite fluoride is synthesized using ammonium fluoride (NH<sub>4</sub>F) as electroactive material of SC at the first time. Different NH<sub>4</sub>F amounts are used to produce perovskite ZIF67-derived fluorides (ZIF67-N). The optimized ZIF67-N electrode shows specific capacitance (C<sub>p</sub>) of 636.8 F/g at 10 mV/s, owing to small particle size and suitable F to 2-methylimidazole ratio for providing high electronegativity. The ZIF67 and cobalt nickel fluoride prepared using NH<sub>4</sub>F but no 2-methylimidazole (CoNi-N) are synthesized to understand roles of fluorine and 2-methylimidazole on energy storage. The ZIF67 electrode shows much smaller C<sub>p</sub> (1.6 F/g) than ZIF67-N electrode, owing to largely enhanced pore width of ZIF67-N even if surface area is largely reduced when NH<sub>4</sub>F is added during synthesis. The SC comprising optimized ZIF67-N electrodes shows maximum energy density of 27.2 Wh/kg at 650.0 W/kg as well as C<sub>p</sub> retention of 86% and Coulombic efficiency of 100% in 8000 times charge/discharge process.

© 2021 Elsevier B.V. All rights reserved.

1. Introduction

Metal organic framework (MOF) with high surface area and tunable structure has been largely applied on energy storage for recent years [1–4]. Zeolitic imidazolate framework 67 (ZIF67) consisted of cobalt ion center and 2-methylimidazole ligand is one of the potential electroactive materials for supercapacitors (SC) [5–7]. However, the intrinsic nature of ZIF67 is not highly capacitive for storing charges even if ZIF67 possesses high surface area for carrying out large amounts of electrochemical reactions. Numerous ex-situ methods were applied on modifying ZIF67 with high redox activity and electrical conductivity. Zhang and co-workers prepared ZIF-derived carbon using co-carbonization technique and obtained a specific capacitance (C<sub>p</sub>) of 228 F/g at 0.1 A/g [8]. Hu *et al.* assembled SC using ZIF-67@amorphous ZIF electrode and capacity retention of 100% after 2000 cycles was obtained [9]. Zhang *et al.* synthesized amorphous carbon@graphite carbon nanoleaves by carbonization of ZIF-Li(Zn)/ZIF67 nanoleaves and achieved C<sub>p</sub> of 252.1 F/g [10]. Combining ZIF67 with carbon materials is also applied to improve energy storage ability of ZIF67 [11,12]. Jian *et al.* designed cobalt sulfide nanocage derived from ZIF interconnected by carbon nanotubes as electrode material for SC [11]. Sundriya *et al.* synthesized ZIF67 and reduced graphene oxide (rGO) composite using stirring approach and obtained C<sub>p</sub> of 326 F/g at 3 A/g [12].

However, comparing to the ex-situ method, the in-situ method is more likely to reduce the experimental process via directly modifying the process of forming MOF derivatives at the very beginning. Also, the nature of MOF derivatives could be much easier to design using in-situ techniques. It was reported that ligand plays important roles on intrinsic properties of MOF, such as chemical stability, rigidity and flexibility [13–15]. Lv *et al.* proposed that stability of MOF relies on robustness of metal ion/ligand coordination bonds. They demonstrated a ligand-rigidification strategy to enhance stability of MOF, including thirteen Zr-based MOF constructed with Zr<sub>6</sub>O<sub>4</sub>(OH)<sub>4</sub>(-CO<sub>2</sub>)<sub>6</sub> units and corresponding ligands. The rigidifying ligand in ZIF67 to synthesize derivatives may be possible to improve the energy storage ability. Ammonium fluoride has been reported to play

\* Corresponding author at: Department of Chemical Engineering and Biotechnology, National Taipei University of Technology, Taipei, Taiwan.  
E-mail address: [lylin@ntut.edu.tw](mailto:lylin@ntut.edu.tw) (L.-Y. Lin).

<sup>1</sup> The authors are equally contributed.

<https://doi.org/10.1016/j.jalcom.2021.162191>  
0925-8388/© 2021 Elsevier B.V. All rights reserved.

Pin-Yan Lee<sup>a,1</sup> Lu-Yin Lin<sup>a,b,\*,1</sup>

<sup>a</sup> Department of Chemical Engineering and Biotechnology, National Taipei University of Technology, Taipei, Taiwan  
<sup>b</sup> Research Center of Energy Conservation for New Generation of Residential, Commercial, and Industrial Sectors, Taipei, Taiwan

共同作者數 (W3)

無 (x1)

1位通訊作者 (x1)

2位(含)以上通訊作者 (x0.8)

有多位作者 Equal Contribution (x0.8)

\* Corresponding author at: Department of Chemical Engineering and Biotechnology, National Taipei University of Technology, Taipei, Taiwan.  
E-mail address: [lylin@ntut.edu.tw](mailto:lylin@ntut.edu.tw) (L.-Y. Lin).

<sup>1</sup> The authors are equally contributed.

# 【共通】研究彈薪申請表論文面向-B27

查詢W3方式

注意事項:

1. 國際學者的定義:除台灣以外皆是外國，且單位須為學術機構(學校、研究機構)。
2. 需檢附論文第一頁為佐證資料。

(二) 作者排序(W2)：作者排序與相對應的權重。

作者排序	第一作者	通訊作者	第二作者	第三作者	第四作者	第五作者以上
權重 2 (W2)	1	1	0.8	0.6	0.4	0.2

(三) 共同作者數(W3)：

1. 若為通訊作者，且該篇文章有兩位以上通訊作者，則該篇須乘以0.8，惟如與國際學者合著者，該國際學者不受此限。
2. 如申請者同屬該篇文章多位 Equal Contribution 之一，則該篇應乘以0.8。

通訊作者數	1 位通訊作者	2 位(含)以上通訊作者 (若其他通訊作者皆為國際 學者則不需 x0.8)	有多位 Equal Contribution
權重 3 (W3)	1	0.8	0.8

3. W2非為通訊作者者，W3請直接勾選【無】；  
W2為通訊作者者，W3請依實際狀況勾選【1位通訊作者、  
2位(含)以上通訊作者、有多位作者】。

情境模擬	w2	w3
為第一作者	1	1
為通訊作者，只有一位	1	1
為通訊作者，且兩位以上 (都是國內學者)	1	0.8
為通訊作者，且兩位以上 (除了申請人以外都是國外學者)	1	1
第二作者後，且不是通訊作者	對應 點數	1
相同貢獻之第一作者	1	0.8
相同貢獻之通訊作者	1	0.8

# 【共通】研究彈薪申請表論文面向-B28

查詢W4方式-企業

依ppt第30.31頁方式下載論文檔案，作者下方之區域，可以看到企業以莊賀喬教授之論文為例：

本篇文章有企業，對應法規應x1.1



Contents lists available at [ScienceDirect](#)

## Food Chemistry

journal homepage: [www.elsevier.com/locate/foodchem](http://www.elsevier.com/locate/foodchem)

### 3D-flower-like porous neodymium molybdate nanostructure for trace level detection of organophosphorus pesticide in food samples

Muthusankar Ganesan<sup>a,c</sup>, Ramadhass Keerthika Devi<sup>b</sup>, Ai-Ho Liao<sup>d,e</sup>, Kuo-Yu Lee<sup>f</sup>, Gopu Gopalakrishnan<sup>c</sup>, Ho-Chiao Chuang<sup>a,\*</sup>

<sup>a</sup> Department of Mechanical Engineering, National Taipei University of Technology, Taipei, 10608, Taiwan, ROC  
<sup>b</sup> Department of Chemical Engineering and Biotechnology, College of Engineering, National Taipei University of Technology, Taipei 106, Taiwan, ROC  
<sup>c</sup> Department of Industrial Chemistry, Alagappa University, Karaikudi 630003, Tamil Nadu, India  
<sup>d</sup> Graduate Institute of Biomedical Engineering, National Taiwan University of Science and Technology, Taipei, Taiwan  
<sup>e</sup> Department of Biomedical Engineering, National Defense Medical Center, Taipei, Taiwan  
<sup>f</sup> SV Probe Technology Taiwan **Co., Ltd.** Zhubei, Taiwan

額外加權 (W4)

- 無 (x1)
- 企業 (x1.1)
- SDG (x1.1)
- SSCI (x1.5)
- 企業、SDG (x1.2)
- 企業、SSCI (x1.6)
- SDG、SSCI (x1.6)
- 企業、SDG、SSCI (x1.8)

# 【共通】研究彈薪申請表論文面向-B29


查詢W4方式-企業

企業通常為 **crop**、**Ltd**、**醫院**，若非前述情況，可於Scival上查詢是否屬企業

Step1: 登入Scival，點選左列【房子圖案】，輸入欲查詢企業(以工研院Industrial Technology Research Institute為例)後，點選欲查詢企業

Step2: 點選【More details on this Institution】，即顯示出欲查詢企業之類型

\*工研院查詢結果為政府機構，故不能勾選企業



# 【共通】研究彈薪申請表論文面向-B30

查詢W4方式-SDG(方法一)

Step1:依ppt第18~21頁方式查詢論文，帶入論文資料畫面後，點選左下方標籤【永續發展目標2023】

Step2:頁面將自動跳轉至此，將隱藏欄位點出即可看見論文所屬之SDG類別

Step3:確定有SDG後，即可勾選對應欄位，並請檢附查詢畫面當作佐證資料

被 0 篇文獻引用  
當本文獻在 Scopus 中被引用時通知我:  
設定引用新通知 >

相關文獻  
Development of thermosensitive poloxamer 407-based microbubble gel with ultrasound mediation for inner ear drug delivery  
Liao, A.-H., Shih, C.-P., Li, M.-W. (2021) Drug Delivery  
Effects of microbubble size on ultrasound-

摘要  
作者關鍵字  
已索引的關鍵字  
永續發展目標 2023  
裝置商品名  
熱門主題  
化學品和 CAS 登記編號  
計量

摘要  
We have previously applied ultrasound (US) with microbubbles (MBs) to enhance drug delivery, with most experiments conducted using single-frequency, high-power density treatments. In the present study, the treatment efficacy was enhanced and concerns were addressed using a combination of low-power-density, single-transducer, dual-frequency US ( $I_{SPTA} = 213 \text{ mW/cm}^2$ ) and MBs of different sizes coated with insulin-like growth factor 1 (IGF-1). This study is the first to investigate the drug-coating capacity of human serum albumin (HSA) MBs of different particle sizes and their drug delivery efficiency. The concentration of HSA was adjusted to produce different MB sizes. The drug-coating efficiency was significantly higher for large-sized MBs than for smaller MBs. In vitro Franz diffusion experiments showed that the combination of dual-frequency US and large MB size delivered the most IGF-1 ( $24.3 \pm 0.47 \text{ ng/cm}^2$ ) to the receptor side at the second hour of treatment. In an in vivo

以莊賀喬教授之論文為例:

額外加權 (W4)

- 無 (x1)
- 企業 (x1.1)
- SDG (x1.1)
- SSCI (x1.5)
- 企業、SDG (x1.2)
- 企業、SSCI (x1.6)
- SDG、SSCI (x1.6)
- 企業、SDG、SSCI (x1.8)



# 【共通】研究彈薪申請表論文面向-B31

查詢W4方式-SDG(方法二)

Step1:登錄至SciVal，輸入老師名字後，點選【View list of publications】

Step2:篩選所欲查詢教師之機構、年份區間與文獻類型後，點選【Apply filter】

(本範例篩選條件為:北科大、2022、Article or Review)

Step3:等Apply filter按鈕反灰後，點選【Export spreadsheet】

以陳生明教授為例:

The screenshot displays the SciVal profile for Chen, Shenming. The overall research performance section shows 946 Scholarly Output, 1.66 Field-Weighted Citation Impact, and 27,988 Citation Count. A red box highlights the 'View list of publications' link. The publications list shows 114 publications, with filters applied for Author (Chen, S.-M.), Year (2022), and Document Type (Article or Review). A red box highlights the 'Export spreadsheet' button in the top right corner of the publications list.

Title	Authors	Year	Scopus Source	
Development of Palladium on Bismuth Sulfide Nanorods as a Bifunctional Nanomaterial for Efficient Electrochemical Detection and Photoreduction of Hg(II) Ions	Veerakumar, P., Jaysiva, G., Chen, S.-M. and 1 more	2022	ACS Applied Materials and Interfaces	
Tailored construction of one-dimensional TiO <sub>2</sub> /Au nanofibers: Validation of an analytical assay for detection of diphenylamine in food samples	Kokulnathan, T., Vishnuraj, R., Chen, S.-M. and 5 more	2022	Food Chemistry	31
UV light assisted photocatalytic degradation of textile waste water by Mg <sub>0.8</sub> -xZnxFe <sub>2</sub> O <sub>4</sub> synthesized by combustion method and in-vitro antimicrobial activities	Bessy, T.C., Bindhu, M.R., Johnson, J. and 3 more	2022	Environmental Research	29
High-performance electrochemical sensing of hazardous pesticide Paraoxon using BiVO <sub>4</sub> nano dendrites equipped catalytic strips	Gopi, P.K., Ngo, D.B., Chen, S.-M. and 2 more	2022	Chemosphere	24
Fabrication of thulium metal-organic frameworks based smartphone sensor towards arsenical feed additive drug detection: Applicable in food safety analysis	Chinnapaiyan, S., Rajaji, U., Chen, S.-M. and 3 more	2022	Electrochimica Acta	23
A robust combination of dysprosium vanadate/halloysite nanotubes: the electrochemical system for dimetridazole detection	Kokulnathan, T., Chen, T.-W., Chen, S.-M. and 4 more	2022	Materials Today Chemistry	21

# 【共通】研究彈薪申請表論文面向-B32

查詢W4方式-SDG(方法二)

Step4:勾選欲匯出之項目再點選【Export CSV】或【Export XLSX】，即可下載檔案

**Export publications** ✕

Select the fields you want to include in the export for your selected publications. Last selected options are remembered. \* in publication year

Select all | Deselect all | Reset to default selection

Publication basics	Publication details	Author/Affiliations	Publication metrics	Scopus Source related	Topic related
<input checked="" type="checkbox"/> Title	<input type="checkbox"/> Reference	<input type="checkbox"/> Scopus Affiliation IDs	<input type="checkbox"/> Views	<input type="checkbox"/> Volume	<input type="checkbox"/> Topic Cluster name
<input checked="" type="checkbox"/> Authors	<input type="checkbox"/> Abstract	<input type="checkbox"/> Scopus Affiliation names	<input type="checkbox"/> Field-Weighted Views Impact	<input type="checkbox"/> Issue	<input type="checkbox"/> Topic Cluster number
<input checked="" type="checkbox"/> Year	<input type="checkbox"/> EID (Scopus ID)	<input type="checkbox"/> Number of Authors	<input type="checkbox"/> Citations	<input type="checkbox"/> Pages	<input type="checkbox"/> Topic name
<input checked="" type="checkbox"/> Full date	<input type="checkbox"/> PubMed ID	<input type="checkbox"/> Scopus Author IDs	<input type="checkbox"/> Field-Weighted Citation Impact	<input type="checkbox"/> Article number	<input type="checkbox"/> Topic number
<input type="checkbox"/> Scopus Source title	<input checked="" type="checkbox"/> Sustainable Development Goals (2023)	<input type="checkbox"/> Scopus Author ID First Author	<input type="checkbox"/> Field-Citation Average	<input type="checkbox"/> ISSN	<input type="checkbox"/> Topic Cluster Prominence Percentile
<input checked="" type="checkbox"/> DOI	<input type="checkbox"/> All Science Journal Classification (ASJC)	<input type="checkbox"/> Scopus Author ID Last Author	<input type="checkbox"/> Outputs in Top Citation Percentiles, per percentile	<input type="checkbox"/> Source ID	<input type="checkbox"/> Topic Prominence Percentile
<input type="checkbox"/> Publication type	<input type="checkbox"/> Code	<input type="checkbox"/> Scopus Author ID Corresponding Author	<input type="checkbox"/> Field-Weighted Outputs in Top Citation Percentiles, per percentile	<input type="checkbox"/> Source type	
<input type="checkbox"/> Open Access	<input type="checkbox"/> Field name	<input type="checkbox"/> Scopus Author ID Single Author	<input type="checkbox"/> Patent citations	<input type="checkbox"/> CiteScore*	
<input type="checkbox"/> Institutions	<input type="checkbox"/> Quacquarelli Symonds (QS)	<input type="checkbox"/> Country/Region		<input type="checkbox"/> CiteScore percentile*	
<input type="checkbox"/> Number of Institutions	<input type="checkbox"/> Code			<input type="checkbox"/> SNIP*	
<input type="checkbox"/> Language	<input type="checkbox"/> Field name			<input type="checkbox"/> SNIP percentile*	
	<input type="checkbox"/> Times Higher Education (THE)			<input type="checkbox"/> SJR*	
	<input type="checkbox"/> Code			<input type="checkbox"/> SJR percentile*	
	<input type="checkbox"/> Field name				
	<input type="checkbox"/> ANZSRC FoR (2020)				
	<input type="checkbox"/> Code				
	<input type="checkbox"/> Field name				

50

Cancel **Export CSV** **Export XLSX**

續下頁

# 【共通】研究彈薪申請表論文面向-B33

查詢W4方式-SDG(方法二)

Step5:匯出的表單即會列出被收錄SDG之論文，確定有SDG後，即可勾選對應欄位，並請檢附匯出表單當作佐證資料

Data set	Publications of Chen, Shenming					
Year range	2013 to 2022					
Subject cla	ASJC					
Filtered by	not filtered					
Types of p	All publication types					
Self-citatic	-					
Data sourc	Scopus					
Date last u	1 November 2023					
Date expor	8 November 2023					
10 publications match the selected filter options:						
Authors	Chen, Shenming -m AND Akilarasan, Muthumariappan					
Publication	2022					
Publication	(Review OR Article)					
Title	Authors	Year	Scopus Source title	DOI	Publication type	Sustainable Development Goals (2023)
Tailored ar	Keerthika I	2022	Food Chemistry	10.1016/j.foodchem.2022.133791	Article	-
In-situ con	Nataraj, N.	2022	Chemical Engineering Journal	10.1016/j.ccej.2022.137025	Article	-
Se substitu	Nataraj, N.	2022	Chemosphere	10.1016/j.chemosphere.2022.134765	Article	SDG 3
Rational sy	Akilarasan	2022	Bioelectrochemistry	10.1016/j.bioelechem.2022.108145	Article	-
Electroche	Yamuna, A	2022	Journal of Electroanalytical Chemistry	10.1016/j.jelechem.2021.115978	Article	-
Fabrication	Sundaresar	2022	Micromachines	10.3390/mi13060876	Article	-
One-pot sy	Babulal, S.	2022	Materials Today Chemistry	10.1016/j.mtchem.2022.101132	Article	SDG 3
Rationally	Tamilalaga	2022	Colloids and Surfaces A: Physicochemical and Engineering Aspects	10.1016/j.colsurfa.2022.129941	Article	-
Synthesis c	Maheshwa	2022	Bioelectrochemistry	10.1016/j.bioelechem.2022.108166	Article	-
In-situ syn	Akilarasan	2022	Process Safety and Environmental Protection	10.1016/j.psep.2022.07.011	Article	-

額外加權  
(W4)

- 無 (x1)
- 企業 (x1.1)
- SDG (x1.1)
- SSCI (x1.5)
- 企業、SDG (x1.2)
- 企業、SSCI (x1.6)
- SDG、SSCI (x1.6)
- 企業、SDG、SSCI (x1.8)

# 【共通】研究彈薪申請表論文面向-B34

查詢W4方式-SSCI (方法一)

Step1:依ppt第18~21頁方式查詢論文，帶入論文資料畫面後，點選【全文選項】

Step2:顯示出下拉選單後，點選【SCIE】，直接帶出Wos查詢畫面

Step3:確定有SSCI後，即可勾選對應欄位，並請檢附查詢畫面當作佐證資料

Interactive Learning Environments • 2021

Potential effects of a role-playing digital gaming learning system on the learning performance and motivation in a humanities course

Chin, Kai-Yi<sup>a</sup> ; Chen, Yen-Lin<sup>b</sup>

45 瀏覽次數

全文選項

以陳彥霖教授之論文為例：

View at Publisher

NTUT Full-Text

Full Text

SCIE

SSCI

Interlibrary Loan

INTERACTIVE LEARNING ENVIRONMENTS

Publisher: ROUTLEDGE JOURNALS, TAYLOR & FRANCIS LTD, 2-4 PARK SQUARE, MILTON PAR

ISSN / eISSN: 1049-4820 / 1744-5191

Web of Science Core Collection: Social Sciences Citation Index (SSCI)

Additional Web of Science Indexes: Current Contents Social And Behavioral Sciences | Essential Science Indicators

額外加權 (W4)

- 無 (x1)
- 企業 (x1.1)
- SDG (x1.1)
- SSCI (x1.5)
- 企業、SDG (x1.2)
- 企業、SSCI (x1.6)
- SDG、SSCI (x1.6)
- 企業、SDG、SSCI (x1.8)

# 【共通】研究彈薪申請表論文面向-B35

查詢W4方式-SSCI (方法二)

Step1:依ppt第18~21頁方式查詢論文，帶入論文資料畫面後，點選期刊名稱，視窗右邊即顯示出來源出版物詳情預覽欄位，點選【瀏覽完整的來源出版物詳情】

Step2:點選【SCIE】，直接帶出Wos查詢畫面

Step3:確定有SSCI後，即可勾選對應欄位，並請檢附查詢畫面當作佐證資料

以陳彥霖教授之論文為例：

The screenshot shows a research article page for "Interactive Learning Environments" (2021) by Chin, Kai-Yi and Chen, Yen-Lin. The article title is "Potential effects of a role-play learning system on the learning and motivation in a human-computer interaction".

Annotations with red boxes and arrows indicate the following steps:

- Step 1: Click on the journal name "Interactive Learning Environments" in the article title area.
- Step 2: Click on the link "瀏覽完整的來源出版物詳情" (View complete source publication details) in the preview window.
- Step 3: Click on the "SCIE" button in the "Library Catalogue" dropdown menu.

The "來源出版物詳情" (Source Publication Details) window shows the following information:

- Journal: Interactive Learning Environments
- Scopus 涵蓋年度: 1990, 從 1992 到 1994, 從 2004 至今
- 圖書館訂閱: 從 January 1998 到 December 2005
- 發表者: Taylor & Francis
- 國際標準期刊號: 1049-4820
- 學科類別: Social Sciences: Education, Computer Science: Computer Science Applications
- 來源出版物種類: 期刊

The "CiteScore 排名" table shows the following data:

ASJC 類別	四分位數	百分位數	數
Education	Q1	98th	24 / 1469
Computer Science Applications	Q1	92nd	57 / 792

The "Library Catalogue" dropdown menu shows the "SCIE" button selected.

The "INTERACTIVE LEARNING ENVIRONMENTS" journal information shows:

- Publisher: ROUTLEDGE JOURNALS, TAYLOR & FRANCIS, 2-4 PARK SQUARE, MILTON PARK, ABINGDON, ENGLAND, OXON, OX14 4RN
- ISSN / eISSN: 1049-4820 / 1744-5191
- Web of Science Core Collection: Social Sciences Citation Index (SSCI)
- Additional Web of Science Indexes: Current Contents Social And Behavioral Sciences | Essential Science Indicators

# 【共通】研究彈薪申請表論文面向-B36

查詢W4方式-SSCI (方法三)

Step1:依ppt第18~21頁方式查詢論文，帶入論文資料畫面後，複製期刊名稱

Step2:至Wos將期刊名稱貼上後，點選【搜尋】，直接帶出查詢畫面

Step3:確定有SSCI後，即可勾選對應欄位，並請檢附查詢畫面當作佐證資料

以陳彥霖教授之論文為例:

The screenshot shows a search result for the journal 'Interactive Learning Environments'. A red box highlights the journal name in the search bar, and another red box highlights the SSCI category 'EDUCATION & EDUCATIONAL RESEARCH - SSCI' in the journal information panel. A red arrow points from the journal name to the SSCI category.

**Interactive Learning Environments**

Potential effects of a role-playing learning system on the learning performance and motivation in a humanities course

**INTERACTIVE LEARNING ENVIRONMENTS**

ISSN: 1049-4820  
EISSN: 1744-5191  
JCR ABBREVIATION: INTERACT LEARN ENVIR  
ISO ABBREVIATION: Interact. Learn. Environ.

**Journal information**

CITATION: Social Sciences Citation Index (SSCI)

CATEGORY: **EDUCATION & EDUCATIONAL RESEARCH - SSCI (SSCI)**

LANGUAGES: English  
REGION: ENGLAND  
1ST ELECTRONIC JCR YEAR: 2005

**Publisher information**

PUBLISHER: ROUTLEDGE JOURNALS, TAYLOR & FRANCIS LTD  
ADDRESS: 2-4 PARK SQUARE, MILTON PARK, ABINGDON OX14 4RN, OXON, ENGLAND  
PUBLICATION FREQUENCY: 8 issues/year

# 【共通】研究彈薪申請表論文面向-B37

查詢W4方式

注意事項:

1. 企業的定義:crop、Ltd、醫院，或Scival上認列之企業。
2. 需檢附論文第一頁為佐證資料，勾選SDG或SSCI者，請檢附查詢畫面為佐證資料。

(四) 額外加權(W4)：若該篇文章與下列合著之加權相對應權重如下所示，有多項加權者請選擇相對應之選項。

額外加權	無	企業	SDG	SSCI
權重4(W4)	1	1.1	1.1	1.5

註一：符合多項加權時，請依表格填寫。

額外加權  
(W4)

- 無(x1)
- 企業 (x1.1)
- SDG (x1.1)
- SSCI (x1.5)
- 企業、SDG (x1.2)
- 企業、SSCI (x1.6)
- SDG、SSCI (x1.6)
- 企業、SDG、SSCI (x1.8)

# 【共通】研究彈薪申請表論文面向-B38

查詢W5方式

依ppt第30.31頁方式下載論文檔案，作者下方之區域，可以看到國際學者以陳生明教授之論文為例：

本篇文章與3位國際學者合著，對應法規應x1.2

Disposable cerium oxide/graphene nanosheets based sensor for monitoring acebutolol in environmental samples and bio-fluids

Subash Vetri Selvi <sup>a,1</sup>, Nandini Nataraj <sup>a,1</sup>, Tse-Wei Chen <sup>a,b,c</sup>, Shen-Ming Chen <sup>a,\*</sup>, Prakash Balu <sup>e</sup>, Xiaoheng Liu <sup>d,\*</sup>

<sup>a</sup> Electroanalysis and Bioelectrochemistry Lab, Department of Chemical Engineering and Biotechnology, National Taipei University of Technology, No. 1, Section 3, Chung-Hsiao East Road, Taipei 106, Taiwan, ROC

<sup>b</sup> Research and Development Center for Smart Textile Technology, National Taipei University of Technology, No.1, Section 3, Chung-Hsiao East Road, Taipei 106, Taiwan

<sup>c</sup> Department of Materials, Imperial College London, London SW7 2AZ, United Kingdom

<sup>d</sup> Key Laboratory of Education Ministry for Soft Chemistry and Functional Materials, Nanjing University of Science and Technology, Nanjing 210094, China

<sup>e</sup> Department of Biotechnology, School of Life Science, Vels Institute of Science, Technology and Advanced Studies, Chennai, Tamilnadu, India

國際合著學術  
機構國家數  
(W5)

- 無 (x1)
- 1-2個國家 (x1.1)
- 3個國家以上 (x1.2)

注意事項：

國際學者的定義：除台灣以外皆是外國，且單位須為學術機構(學校、研究機構)。



# 【共通】研究彈薪申請表論文面向-B39

查詢W5方式-國際學者

國際學者通常為 **University**、**Academic**、**College**、**Laboratory**，若非前述情況，可於 Scival 上查詢是否屬研究機構，查詢方式同前

\*私人公司之研究室不屬於研究機構

The screenshot shows the Scival interface for the National Taipei University of Technology. The left sidebar has a red box around the 'Institutions and Groups' icon. Below it, the 'National Taipei University of Technology' is listed in the 'Favorites' section, also with a red box. A red arrow points from the 'More details on this Institution' link to a pop-up window. In this window, the 'Institution type: Academic' is highlighted with a red box. Below this, there are options for 'About', 'Compare definitions', and 'Groups'. The 'Institution definition' section includes a checkbox for 'Unknown' and a button to 'Create Research Area with listed affiliations'.

# 【共通】研究彈薪申請表注意事項

## 溫馨小提醒

1	請勿使用舊版、未更新為現行查詢結果或非紙本出版年度之佐證	9	論文未被Scopus或WOS收錄，或論文發表當年度期刊未被Scopus或WOS收錄，將不予採計點數
2	請確證論文機構是否被列為China、PROC等不符合教育部列名原則之情事，如有則不予計點	10	使用之論文應符合申請區間 (請以紙本出刊為主) 例：112年度申請時可用論文為2018~2022，故2023發表之論文則不可納入計點
3	學術論著績效僅限 <b>Original article</b> 與 <b>Review article</b> 兩類期刊論文點數， <b>研討會論文及其他類型文章均不予計點</b>	11	請使用正確/最新版之表格，並勿更動研發處查詢之本校FWCI與h5值
4	Ranking值請計算至小數點後2位，勿四捨五入	12	多年期國科會計畫應分年度計算經費與點數，並請提供教評系統列印文件作為佐證
5	同篇文章有2位以上通訊作者，則該篇須乘上0.8 如與國際學者合著者，則該國際學者不受此限(不需乘0.8)	13	國科會計畫點數依學院慣例審查，不須送至研發處蓋章
6	如有多位(含2位)以上國內通訊作者，第一位通訊作者亦應乘0.8	14	國科會產學合作案應列入產學，不計入專題計畫點數
7	企業的定義:crop、Ltd、醫院 國際學者的定義:除台灣以外皆是外國，且單位須為學術機構(學校、研究機構)	15	產學合作與技轉點數應先送產學處審核確認，檢核後即可送交系辦彙整，不須先送至研發處
8	同篇文章有2位以上本校教師為共同作者，請檢附其他教師同意書	16	A類申請項目可複選，惟額外項目不得單獨勾選且皆須檢附佐證